

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

( Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Lohit**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May, 2017
Rainfall =72.2 mm (Tezu)	Partly cloudy weather with possibility of moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 20-22°C and 14-15°C respectively. Expected morning and evening relative humidity will be 81-97% and 48-58% respectively. Wind direction is E with speed range of 4-5 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• Rice: Spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC @ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>• Provide sugar syrup near the box</li> <li>• Add insulation and ventilation systems</li> </ul>
<b>LIVESTOCKS</b>			

<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Lower Dibang Valley**

No: 38/2017/Fri

Dated: 12<sup>th</sup> May, 2017



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May, 2017
Rainfall = 32.3 mm (Roing)	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 24-27°C and 9-10°C respectively. Expected morning and evening relative humidity will be 80-97% and 36-50% respectively. Wind direction is mainly E with speed range of 3-5 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• Carry out thinning and earthing up in maize without delay</li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing to avoid fungal</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• Denavelling (removal of male bud) to improve bunch weight</li> <li>• Bagging of bunches to protect against unusual rainfall.</li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilled bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>• Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</li> </ul>
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Citrus	Flowering / Fruiting	Bark eating caterpillar	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>• Budding method of propagation (T-budding) to commence in oranges using Rangpur lime as rootstock.</li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• Spray (Planofix) NAA based growth regulator @2m/5lit water during flowering stage in orange plant to reduce fruit/flower drop</li> <li>• For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</li> <li>• Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			

Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>Box must be covered with straw/dry leaves to provide warmth</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>Animals should be given lukewarm feed and water.</li> <li>Keep the houses dry and clean.</li> <li>1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>Avoid overcrowding.</li> <li>Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>Remove wet litter/provide artificial heating and light</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>Separate sick animals</li> <li>The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Lower Subansiri**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Weather summary of the previous week	Weather forecast valid up to 14 <sup>th</sup> May 2017
Rainfall = 0.0 mm (Ziro)	Mostly cloudy with possibility of light/moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 25-28°C and 14-15°C respectively. Expected morning and evening relative humidity will be 88-93% and 48-75% respectively. Wind direction is variable with speed range of 2-4 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Pulses	Vegetative		<ul style="list-style-type: none"> <li>• <b>Apply biosulphur @2ml/lit water to enhance yield</b></li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
<i>Jhum</i> field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>• <b>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</b></li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> <li>• On clear/dry weather or using sticker (@1ml/lit), spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> </ul>

			<ul style="list-style-type: none"> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>• Box must be covered with straw/dry leaves to provide warmth</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Papum Pare**

No: 38/2017/Fri

Dated: 12<sup>th</sup> May, 2017



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall: 0.8 mm; T. Max: 35.0°C; T. Min: 20.0°C ; RH: 64-82%	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 25-33°C and 20-21°C respectively. Expected morning and evening relative humidity will be 80-96% and 40-80% respectively. Expected wind direction is variable with speed range of 2-4 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Pulses	Vegetative		<ul style="list-style-type: none"> <li>• <b>Apply biosulphur @2ml/lit water to enhance yield</b></li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> <li>• On clear/dry weather or using sticker (@1ml/lit), spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• <b>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</b></li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain</li> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width,</li> </ul>

			<p>15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</p> <ul style="list-style-type: none"> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>• Box must be covered with straw/dry leaves to provide warmth</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Tawang**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall = 9.0 mm	Mostly cloudy with possibility of light/moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 16-18°C and 6-7°C respectively. Expected morning and evening relative humidity will be 83-89% and 52-68% respectively. Wind direction is mostly ENE & E with probable speed range of 4-7 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- All spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>Carry out thinning and earthing up in maize without delay</li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize</li> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> <li>On clear/dry weather or using sticker (@1ml/lit), spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li><b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li><b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <u>25EC@0.05%</u> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain</li> <li>Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li><b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li><b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li><b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>Provide sugar syrup near the box</li> <li>Add insulation and ventilation systems</li> </ul>

## LIVESTOCKS

Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds.</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>

**Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247**

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Tirap**

No: 38/2017/Fri

Dated: 12<sup>th</sup> May, 2017



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May, 2017
Rainfall = 0.8 mm (Deomali)	<b>Variable weather with possibility of light rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 27-33°C and 22-23°C respectively. Expected morning and evening relative humidity will be 81-96% and 46-83% respectively. Wind direction is mainly SSE with speed range of 2-4 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated.**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• Carry out thinning and earthing up in maize without delay</li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> <li>• Careful timing of planting can avoid damage; once the tillering stage is over, attack of some insects reduces.</li> <li>• Follow clean cultivation, presence of grasses near the rice crop favors build-up of pest population</li> <li>• Seedling root dip treatment in 0.02% of Chlorpyrifos 20EC for 12 hours before transplanting protects the crop for 20 to 30 days</li> <li>• Avoid closer spacing while transplantation which creates favorable microclimate for pests in the field. Follow wider spacing of 20X15cm.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>• Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• On clear day, spray Azadirachtin 0.15EC @ 6ml/lit water against insect pests</li> <li>• In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• Denavelling (removal of male bud) to improve bunch weight</li> <li>• Bagging of bunches to protect against unusual rainfall.</li> </ul>
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Citrus	Flowering/fruiting	Bark eating caterpillar	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole 25EC@0.05% against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain</li> <li>• For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</li> <li>• Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm</li> </ul>

			<p>between rows and 25cm between plants.</p> <ul style="list-style-type: none"> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
			<ul style="list-style-type: none"> <li>• Box must be covered with straw/dry leaves to provide warmth</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Upper Siang**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May, 2017
Rainfall = 0.0 mm (Tuting)	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 29-31°C and 20-21°C respectively. Expected morning and evening relative humidity will be 71-85% and 41-52% respectively. Wind direction is mainly SSE & S with speed range of 3-5 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- Disinfecting the animal/bird sheds with lime powder/bleaching powder/4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost poly-houses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• On clear/dry weather or using sticker (@1ml/lit), spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
<i>Jhum</i> field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>• <b>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</b></li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <u>25EC@0.05%</u> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain</li> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm</li> </ul>

			<p>between rows and 25cm between plants.</p> <ul style="list-style-type: none"> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>• Box must be covered with straw/dry leaves to provide warmth</li> <li>• Provide sugar syrup</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide broad-spectrum antihelminthic drugs under vet supervision and recommended doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR. BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

( Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




भा.क.अनु.प. - पू.प.सं.  
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Upper Subansiri**

No: 38/2017/Fri

Dated: 12<sup>th</sup> May, 2017



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall = 0.0 mm (Puchi Geko); Daporjo: 0.0 mm	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 25-28°C and 14-15°C respectively. Expected morning and evening relative humidity will be 88-92% and 53-64% respectively. Expected wind direction is mainly SW & WSW with speed range of 2-4 kmph.
<b>Warning</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Avoid spray of insecticides during flowering stage as it may hinder pollination by honey bees and other insects.
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li><b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li><b>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</b></li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li><b>On clear day, spray Azadirachtin 0.15EC @ 6ml/lit water against insect pests</b></li> <li><b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li><b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li><b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <u>25EC@0.05%</u> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain</li> <li><b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li><b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li><b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> </ul>
Apple	Vegetative	Wooly aphids	<ul style="list-style-type: none"> <li>Swab the stem with mineral oil against wooly aphid.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			

Honey bees			<ul style="list-style-type: none"> <li>• Provide sugar syrup near the box</li> <li>• Add insulation and ventilation systems</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Provide adequate bedding material (paddy straw, rice husk) to newly born piglets.</li> <li>• Animals should be given lukewarm feed and water.</li> <li>• All animals should be tethered in a covered shelter. Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Provide broad-spectrum antihelminthic drugs under vet supervision and recommended doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit**  
**ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: West Kameng**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall = Bomdila: 0.0 mm; Bhalukpong: 10.4 mm	<b>Partly cloudy with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 17-21°C and 10-11°C respectively. Expected morning and evening relative humidity will be 79-91% and 48-72% respectively. Wind direction is mainly NNE & ENE with speed range of 3-5 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**
- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>Rice: Spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li><b>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</b></li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li><b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li><b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <u>25EC@0.05%</u> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain</li> <li>Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li><b>remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li><b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Banana	Vegetative		<ul style="list-style-type: none"> <li>Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li><b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li><b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li><b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> </ul>

Apple	Vegetative	Woolly aphids	<ul style="list-style-type: none"> <li>Swab the stem with mineral oil against woolly aphid.</li> </ul>
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>Provide sugar syrup near the box</li> <li>Add insulation and ventilation systems</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>Separate sick animals</li> <li>The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>Animals should be given lukewarm feed and water.</li> <li>Keep the houses dry and clean.</li> <li>1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>Avoid overcrowding.</li> <li>Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका Agro-met Advisory Bulletin

( Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: West Siang**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Basar: Latitude: 27°59.53/Longitude: 94.41.269/Altitude: 616 amsl

Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall = 6.8 mm. T. Max: 30.5°C, T. Min: 15.2°C, RH: 51-92%, Wind Speed: 5-6 kmph, Wind Direction: NE & WN	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 29-32°C and 20-22°C respectively. Expected morning and evening relative humidity will be 86-92% and 48-66% respectively. Wind direction is mainly W & WNW with speed range of 2-4 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

## Weather Based Agro-meteorological Advisories Bulletin

### General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li><b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Pulses	Vegetative		<ul style="list-style-type: none"> <li><b>Apply biosulphur @2ml/lit water to enhance yield</b></li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li><b>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</b></li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li><b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li><b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li><b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li><b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li><b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scab in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li><b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li><b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> </ul>
<b>Honey Bees</b>			

Honey bees			<ul style="list-style-type: none"> <li>• Box must be covered with straw/dry leaves to provide warmth</li> <li>• Provide sugar syrup</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		<ul style="list-style-type: none"> <li>• Filling of nursery polybags for raising of takopalm seedlings</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Provide adequate bedding material (paddy straw, rice husk) to newly born piglets.</li> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Anjaw**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Hawai: Latitude: 28.2179994/ Longitude: 94.7277528/Altitude: 1296 m amsl

Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
No weather summary available	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 30-31°C and 21-24°C respectively. Expected morning and evening relative humidity will be 81-94% and 38-46% respectively. Wind will blow mostly E & ESE at a speed range of 4-6 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **Warning: Spraying of DDT in crops is lethal**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li><b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
<i>Jhum</i> field	Sowing		<ul style="list-style-type: none"> <li>Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> </ul>
Pulses	Vegetative		<ul style="list-style-type: none"> <li><b>Apply biosulphur @2ml/lit water to enhance yield</b></li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li><b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li><b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Orchards	Multiple stages		<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Banana	Vegetative		<ul style="list-style-type: none"> <li>Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li><b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li><b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li><b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/ Citrus scab	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scab in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</li> <li>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</li> <li>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> <li>Provide adequate drainage</li> </ul>
<b>AGROFORESTRY</b>			
Toko palm	Nursery		<ul style="list-style-type: none"> <li>Filling of nursery polybags for raising of takopalm seedlings</li> </ul>

<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision against FMD</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide broad-spectrum antihelminthic drugs under vet supervision and recommended doses.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>• Separate sick animals</li> <li>• The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>• Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. Pregnant animals above 7 months of pregnancy should be avoided from vaccination.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा

**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Changlang**

No: 38/2017/Fri

Dated: 12<sup>th</sup> May, 2017



Changlang (HQ): Latitude: 27.1305856/Longitude: 95.7342373/Altitude: 580 masl

Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May, 2017
Rainfall = Changlang: 0.0 mm Miao: 7.2 mm	<b>Partly cloudy weather forecasted with light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 27-31°C and 20-21°C respectively. Expected morning and evening relative humidity will be 82-99% and 49-68% respectively. Wind direction is mainly E with speed range of 3-4 kmph.
<b>Warning</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Pulses	Vegetative		<ul style="list-style-type: none"> <li>Apply biosulphur @2ml/lit water to enhance yield</li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li>Carry out thinning and earthing up in maize without delay</li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>On clear day, spray Azadirachtin 0.15EC @ 6ml/lit water against insect pests</li> <li>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>Denavelling (removal of male bud) to improve bunch weight</li> <li>Bagging of bunches to protect against unusual rainfall.</li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</li> <li>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</li> <li>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</li> </ul>
Orchards			<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> </ul>
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>Box must be covered with straw/dry leaves to provide warmth</li> <li>Provide sugar syrup</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages	FMD (high fever and appearance	<ul style="list-style-type: none"> <li>Keep the houses dry and clean.</li> <li>1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of</li> </ul>

		of mucous membrane of the mouth and foot.)	age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Dibang Valley**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Anini: Latitude: 28.2179994/Longitude: 94.7277528/Altitude: 1968 masl

Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May, 2017
Rainfall = 0.0 mm (Anini)	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 24-27°C and 9-10°C respectively. Expected morning and evening relative humidity will be 80-91% and 38-54% respectively. Wind direction is mainly E with speed range of 2-5 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Go for spray operation only on dry/clear weather.
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• On clear/dry weather or using sticker (@1ml/lit), spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
<i>Jhum</i> field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>

Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
Apple	Vegetative	Woolly aphids	<ul style="list-style-type: none"> <li>• Swab the stem with mineral oil against woolly aphid.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>• In case of rain provide sugar syrup</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



भा.क.अ.प. - पू.प.सं.  
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

### District: East Kameng

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall = 0.0 mm	<b>Partly cloudy with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 25-28°C and 14-15°C respectively. Expected morning and evening relative humidity will be 82-94% and 47-80% respectively. Expected wind direction is mainly E & NE with speed range of 2-4 kmph.
<b>Warning:</b>	• <b>Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing
- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• On clear/dry weather or using sticker (@1ml/lit), spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> <li>•</li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
<i>Jhum</i> field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>• Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Spray of Difenocomazole <a href="#">25EC@0.05%</a> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>• <b>Mulching ginger beds with locally available</b></li> </ul>

			<b>leaves/materials is advocated to enhance germination</b>
Orchards			<ul style="list-style-type: none"> <li>• Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> <li>• Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>
<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>• Animals should be given lukewarm feed and water.</li> <li>• Keep the houses dry and clean.</li> <li>• 1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>• Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>• Avoid overcrowding.</li> <li>• Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>• Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>• Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>• Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>• 1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>• Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>• Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR. BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: East Siang**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Pasighat: Latitude: 28.0700/Longitude: 95.3300/Altitude: 153 masl

Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2017
Rainfall: 0.0 mm; T. Max: 36.0°C; T. Min: 22.0°C ; RH: 68-86%	<b>Variable weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 28-33°C and 20-22°C respectively. Expected morning and evening relative humidity will be 82-94% and 43-73% respectively. Expected wind direction is mainly W with speed range of 2-4 kmph.
<b>Warning:</b>	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Finger millet	Sowing		<ul style="list-style-type: none"> <li>Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Maize	Knee height		<ul style="list-style-type: none"> <li><b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Pulses	Vegetative		<ul style="list-style-type: none"> <li><b>Apply biosulphur @2ml/lit water to enhance yield</b></li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing to avoid fungal infections during rainy days</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li><b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li><b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li><b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li><b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li><b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>Spray of Difenocomazole <b>25EC@0.05%</b> against citrus scabe in Khasi Mandarin. Mix sticker 1ml/lit in case of rain.</li> <li>Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li><b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li><b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li><b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Orchards			<ul style="list-style-type: none"> <li>Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.</li> </ul>

			<ul style="list-style-type: none"> <li>Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards</li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>The duration of primary nursery being 3-4 months.</li> </ul>
<b>Honey Bees</b>			
Honey bees			<ul style="list-style-type: none"> <li>On prolong rain, provide sugar syrup</li> </ul>
<b>LIVESTOCKS</b>			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>Give broad-spectrum antihelmintic drugs as per prescribed doses.</li> <li>Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit  
ICAR, BASAR**

# कृषि मौसम सलाहकार सेवा पत्रिका

## Agro-met Advisory Bulletin

(Period: 13<sup>th</sup> to 16<sup>th</sup> May 2017)

ग्रामीण कृषि मौसम सेवा



**GRAMIN KRISHI MAUSAM SEWA**  
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region  
Arunachal Pradesh Centre, Basar-791101

**District: Kurung Kumey**

**No: 38/2017/Fri**

**Dated: 12<sup>th</sup> May, 2017**



Koloriang: Latitude: 27.9167°/Longitude: 93.3500°/Altitude: 1040 masl

Weather summary of the previous week	Weather forecast valid up to 17 <sup>th</sup> May 2016
Past weather summary not available	<b>Mostly cloudy weather with possibility of light/moderate rainfall.</b> Daily maximum and minimum temperature for the following period is predicted to be 30-31°C and 20°C respectively. Expected morning and evening relative humidity will be 81-90% and 47-69% respectively. Wind direction is variable with speed range of 2-5 kmph.
Warning	<b>• Rain/thundershowers are likely in few places</b>

### Weather Based Agro-meteorological Advisories Bulletin

#### General Recommendations

- **Spray operations should be done only on dry/clear weather or in extreme case use stickers**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**
- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
<b>FIELD CROPS</b>			
Maize	Knee height		<ul style="list-style-type: none"> <li>• <b>Carry out thinning and earthing up in maize without delay</b></li> </ul>
Finger millet	Sowing		<ul style="list-style-type: none"> <li>• Sowing of short duration finger millet varieties like VL-324 &amp; VR-708 can be carried out without delay.</li> </ul>
Rice	Seedling	Stem borer	<ul style="list-style-type: none"> <li>• Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice and maize.</li> <li>• Rice: Spray Triazophos 40EC @ 4 ml/lit against stem borer in rice nursery as well as in main field.</li> <li>• Cut the tips of the leaves before transplanting reduces the population of stem borer as the eggs are laid on that site. It will reduce the carryover of the infestation to the main field.</li> </ul>
Jhum field	Sowing		<ul style="list-style-type: none"> <li>• Adopt sowing of leguminous crops like cowpea, frenchbean, soybean etc at 2-3 rows across the slope of jhum field at desirable intervals or ratios with rice to prevent soil erosion and conserve moisture and fertility.</li> <li>• Cowpea variety (Kashi Kanchan) and Frenchbean variety (Selection-9) are advocated for intercropping as they are short duration, dwarf and high yielding.</li> <li>• Carry out seed dressing/treatment with Thiram or Carbendazim @ 2g/kg seed before sowing</li> </ul>
Pea	Seedling		<ul style="list-style-type: none"> <li>• Provide proper drainage to already sown crops</li> </ul>
<b>VEGETABLES</b>			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> <li>• Provision of drainage to remove excess water followed by earthing up in vegetables.</li> <li>• On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development.</li> <li>• <b>On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests</b></li> <li>• <b>In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboor + sticks.</b></li> </ul>
<b>HORTICULTURAL CROPS</b>			
Banana	Vegetative		<ul style="list-style-type: none"> <li>• Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance.</li> <li>• <b>Denavelling (removal of male bud) to improve bunch weight</b></li> <li>• <b>Bagging of bunches to protect against unusual rainfall.</b></li> </ul>
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> <li>• Disease free ginger/turmeric rhizomes of 30-40g having atleast 2 eyes to be sown in finely tilth bed of 1 meter width, 15 cm high and of convenient length at a spacing of 30 cm between rows and 25cm between plants.</li> <li>• <b>Mulching ginger beds with locally available leaves/materials is advocated to enhance germination</b></li> </ul>
Citrus	Fruiting	Bark eating caterpillar/	<ul style="list-style-type: none"> <li>• Budding method of propagation (T-budding) to commences in oranges using Rangpur lime as rootstock.</li> <li>• <b>If the citrus trees are showing oozing symptoms of gum then scrap the area with a sharp knife and apply Metalaxyl MZ-72 or Fosetyl A1 paste on it.</b></li> <li>• Cleaning &amp; smearing of trunk base with Bordeaux paste</li> <li>• If flowering begins, avoid spray of insecticides/ pesticides as it may affect pollination process.</li> <li>• <b>For bark eating caterpillar, remove wooden frass and inject Dichlorovos (0.1%) @ 5ml with disposable syringe.</b></li> <li>• <b>Dead wood on the citrus trees should be kept pruned, about 2cm, below the dead portion.</b></li> </ul>
Large Cardamom	Sowing		<ul style="list-style-type: none"> <li>• Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching &amp; regular light irrigation.</li> <li>• The duration of primary nursery being 3-4 months.</li> </ul>

<b>Agroforestry</b>			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
<b>Honey Bees</b>			
			<ul style="list-style-type: none"> <li>Box must be covered with straw/dry leaves to provide warmth</li> </ul>
<b>LIVESTOCKS</b>			
<b>Livestock</b>	<b>Stage</b>	<b>Disease</b>	<b>Precautionary Practices to be followed</b>
Cattle	All stages	FMD	<ul style="list-style-type: none"> <li>1<sup>st</sup> injection at 6-8 weeks of age, 2<sup>nd</sup> injection after 6 months of 1<sup>st</sup> injection followed by annual vaccination under vet supervision</li> <li>Separate sick animals</li> <li>The animal should be washed with lukewarm water added with little potash (KMnO<sub>4</sub>) or neem leaves</li> <li>Long hair near the udder/stomach/back legs should be teamed short.</li> </ul>
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> <li>Provide adequate bedding material (paddy straw, rice husk) to newly born piglets.</li> <li>Animals should be given lukewarm feed and water.</li> <li>All animals should be tethered in a covered shelter. Keep the houses dry and clean.</li> <li>1<sup>st</sup> injection at 6 months of age and 2<sup>nd</sup> injection at 12 months of age followed by annual vaccination under vet supervision</li> </ul>
Poultry	All stages		<ul style="list-style-type: none"> <li>Follow 'All in All out' system of rearing and do not rear birds of different age groups together.</li> <li>Avoid overcrowding.</li> <li>Provide adequate clean drinking water+electrolyte and multi-vitamins</li> <li>Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> <li>➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</li> </ul> </li> <li>Remove wet litter/provide artificial heating and light</li> </ul>
Fishery			<ul style="list-style-type: none"> <li>Best time to prepare the culture pond like cleaning aquatic weeds</li> </ul>
Mithun		FMD	<ul style="list-style-type: none"> <li>Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route.</li> <li>Pregnant animals above 7 months of pregnancy should be avoided.</li> </ul>
<b>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</b>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department  
Contact: - [IAAS.BASAR@gmail.com](mailto:IAAS.BASAR@gmail.com), Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit**  
**ICAR, BASAR**