

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lower Dibang Valley

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June, 2017
Rainfall = 28.9 mm (Roing)	Mostly/partly cloudy with possibility of light to rather heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 18-22°C and 11-13°C respectively. Expected morning and evening relative humidity will be 91-94% and 61-85% respectively. Wind direction is mainly E & ESE with speed range of 3-6 kmph.
Warning:	<ul style="list-style-type: none">• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**
- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Maize	Knee height		<ul style="list-style-type: none"> • Carry out thinning and earthing up in maize without delay
Rice	Nursery/transplanting		<ul style="list-style-type: none"> • Maintain proper drainage for nursery plot of rice. • Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> • Avoid sowing • If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> • Manual weeding advocated • Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Finger millet	Sowing		<ul style="list-style-type: none"> • Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> • Provision of drainage to remove excess water followed by earthing up in vegetables. • In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. • Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting		<ul style="list-style-type: none"> • Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> • Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> • Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. • Denavelling (removal of male bud) to improve bunch weight • Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> • Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. • Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> • Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. • The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> • Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> • Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> • Animals should be given lukewarm feed and water. • Keep the houses dry and clean. • 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> • Give broad-spectrum antihelmintic drugs as per prescribed doses. • Follow 'All in All out' system of rearing and do not rear birds of different age groups together. • Avoid overcrowding. • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet.

			<ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision • Separate sick animals • The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves • Long hair near the udder/stomach/back legs should be teamed short.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lower Subansiri

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th May 2017
Rainfall = NA (Ziro)	Mostly cloudy with possibility of light/moderate rainfall and heavy rainfall on 31 st May. Daily maximum and minimum temperature for the following period is predicted to be 18-22°C and 14-16°C respectively. Expected morning and evening relative humidity will be 96-97% and 78-92% respectively. Wind direction is variable with speed range of 2-3 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Pulses	Vegetative		<ul style="list-style-type: none"> Apply biosulphur @2ml/lit water to enhance yield
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development. On clear day, spray Azadirachtin 0.15EC @ 6ml/lit water against insect pests In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Agroforestry			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision

Poultry	All stages		<ul style="list-style-type: none"> • Give broad-spectrum antihelmintic drugs as per prescribed doses. • Follow 'All in All out' system of rearing and do not rear birds of different age groups together. • Avoid overcrowding. • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision • Separate sick animals • The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves • Long hair near the udder/stomach/back legs should be teamed short.
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प.सं.
ICAR-RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Papum Pare

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
Rainfall: 31.1 mm; T. Max: 36.0°C; T. Min: 21.0°C ; RH: 61-90%	Partly cloudy with possibility of light/moderate rainfall and heavy rainfall on 31st May. Daily maximum and minimum temperature for the following period is predicted to be 20-29°C and 18-22°C respectively. Expected morning and evening relative humidity will be 96-98% and 61-95% respectively. Expected wind direction is mainly E and ENE with speed range of 2-4 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**
- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Pulses	Vegetative		<ul style="list-style-type: none"> Apply biosulphur @2ml/lit water to enhance yield
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development. On clear day, spray Azadirachtin 0.15EC@ 6ml/lit water against insect pests In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision Separate sick animals The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves Long hair near the udder/stomach/back legs should be teamed short.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the	<ul style="list-style-type: none"> 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision

		mouth and foot.)	
Poultry	All stages		<ul style="list-style-type: none"> • Give broad-spectrum antihelmintic drugs as per prescribed doses. • Follow 'All in All out' system of rearing and do not rear birds of different age groups together. • Avoid overcrowding. • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Tawang

No: 42/2017/Fri

Dated: 26th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
Rainfall = 16.0 mm	Mostly cloudy with possibility of light to rather heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 13-16°C and 9-10°C respectively. Expected morning and evening relative humidity will be 86-97% and 65-87% respectively. Wind direction is variable with probable speed range of 2-10 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- All spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. On clear/dry weather, spray Boom (organic liquid manure) @ 4ml/lit of water for proper growth and development. On clear day, spray Azadirachtin 0.15EC @ 6ml/lit water against insect pests In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Provide sugar syrup near the box Add insulation and ventilation systems
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Give broad-spectrum antihelmintic drugs as per prescribed doses. Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-vitamins Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox

			vaccine and 56-70 days: RD R-2B strain.
			<ul style="list-style-type: none"> • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision • Separate sick animals • The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves • Long hair near the udder/stomach/back legs should be teamed short.
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)




भा.क.अनु.प. - पू.प.सं.
ICAR - RCRNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Tirap

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June, 2017
Rainfall = 30.4 mm (Deomali)	Mostly cloudy with possibility of moderate rainfall. Daily maximum and minimum temperature for the following period is predicted to be 23-29°C and 21-23°C respectively. Expected morning and evening relative humidity will be 96-100% and 76-97% respectively. Wind direction is variable with speed range of 2-4 kmph.
Warning:	<ul style="list-style-type: none">Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated.
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Give broad-spectrum antihelmintic drugs as per prescribed doses. Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-vitamins Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days:

			<p>F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.</p> <ul style="list-style-type: none"> • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision • Separate sick animals • The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves • Long hair near the udder/stomach/back legs should be teamed short.
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)



भा क अरु प - पू प सं
ICAR - RCNEH

ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Upper Siang

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June, 2017
Rainfall = 18.2 mm (Tuting)	Variable weather with possibility of light/moderate rainfall and rather heavy rainfall on 31st May. Daily maximum and minimum temperature for the following period is predicted to be 18-22°C and 14-16°C respectively. Expected morning and evening relative humidity will be 79-89% and 62-72% respectively. Wind direction is mainly ESE and SE with speed range of 2-5 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/bleaching powder/4%NaOH at regular interval is advocated
- Spraying operation should be done only on dry/clear weather
- Go for low cost poly-houses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth Provide sugar syrup
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Provide broad-spectrum antihelminthic drugs under vet supervision and recommended doses. Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-vitamins Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adapted fowl pox vaccine and 56-70 days: RD R-2B strain.

			<ul style="list-style-type: none"> Remove wet litter/provide artificial heating and light
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision Separate sick animals The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves Long hair near the udder/stomach/back legs should be teamed short.
Mithun		FMD	<ul style="list-style-type: none"> Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

Agrometeorological Field Unit
ICAR, BASAR

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA

(Integrated Agro-Meteorological Advisory Services)




भा क अरु प - पू प सं
ICAR - RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Upper Subansiri

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
Rainfall = NA (Puchi Geko); Daporijo: 24.0 mm	Mostly cloudy with possibility of light/moderate rainfall and heavy rainfall on 31 st May. Daily maximum and minimum temperature for the following period is predicted to be 15-22°C and 11-13°C respectively. Expected morning and evening relative humidity will be 95-96% and 76-89% respectively. Expected wind direction is variable with speed range of 2-3 kmph.
Warning	Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Avoid spray of insecticides during flowering stage as it may hinder pollination by honey bees and other insects.
- Spraying operation should be done only on dry/clear weather
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Provide sugar syrup near the box Add insulation and ventilation systems
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision Separate sick animals The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves Long hair near the udder/stomach/back legs should be teamed short.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Provide adequate bedding material (paddy straw, rice husk) to newly born piglets. Animals should be given lukewarm feed and water. All animals should be tethered in a covered shelter. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Provide broad-spectrum antihelminthic drugs under vet supervision and recommended doses. Follow 'All in All out' system of rearing and do not rear birds of

			<p>different age groups together.</p> <ul style="list-style-type: none"> • Avoid overcrowding. • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
<p>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</p>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प.सं.
ICAR-RCNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: West Kameng

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
Rainfall = Bomdila: 0.0 mm; Bhalukpong: 6.2 mm	Partly cloudy with possibility of light/moderate rainfall and rather heavy rainfall on 31st May. Daily maximum and minimum temperature for the following period is predicted to be 15-22°C and 11-13°C respectively. Expected morning and evening relative humidity will be 94-98% and 69-92% respectively. Wind direction is variable with speed range of 2-5 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**
- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Apple	Vegetative	Wooly aphids	<ul style="list-style-type: none"> Swab the stem with mineral oil against wooly aphid.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Provide sugar syrup near the box Add insulation and ventilation systems
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision Separate sick animals The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves Long hair near the udder/stomach/back legs should be teamed short.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding.

			<ul style="list-style-type: none"> • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: West Siang

No: 43/2017/Tue

Dated: 30th May, 2017



Basar: Latitude: 27°59.53/Longitude: 94.41.269/Altitude: 616 amsl

Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
<p>Basar: Rainfall = 15.6 mm. T. Max: 31.2°C, T. Min: 17.9°C, RH: 75-92%, Wind Speed: 3-6 kmph, Wind Direction: NE & WN</p> <p>Along: 3.0 mm</p>	<p>Mostly cloudy with possibility of light/moderate rainfall and rather heavy rainfall on 31st May. Daily maximum and minimum temperature for the following period is predicted to be 23-29°C and 21-23°C respectively. Expected morning and evening relative humidity will be 93-96% and 76-89% respectively. Wind direction is variable with speed range of 2-3 kmph.</p>
<p>Warning:</p>	<ul style="list-style-type: none"> Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Pulses	Vegetative		<ul style="list-style-type: none"> Apply biosulphur @2ml/lit water to enhance yield
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth Provide sugar syrup
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision Separate sick animals The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves Long hair near the udder/stomach/back legs should be teamed short.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Provide adequate bedding material (paddy straw, rice husk) to newly born piglets. Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision

Poultry	All stages		<ul style="list-style-type: none"> • Give broad-spectrum antihelmintic drugs as per prescribed doses. • Follow 'All in All out' system of rearing and do not rear birds of different age groups together. • Avoid overcrowding. • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Anjaw

No: 43/2017/Tue

Dated: 30th May, 2017



Hawai: Latitude: 28.2179994/ Longitude: 94.7277528/Altitude: 1296 m amsl

Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
No weather summary available	Partly/mostly cloudy with possibility of light to heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 20-23°C and 15-16°C respectively. Expected morning and evening relative humidity will be 92-96% and 58-80% respectively. Wind direction is mainly E and ESE at a speed range of 3-7 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **Warning: Spraying of DDT in crops is lethal**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Orchards	Multiple stages		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Citrus	Fruiting	Citrus scab/ leaf miner	<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Large Cardamom	Sowing/germination		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months. Provide adequate drainage
Ginger/Turmeric	Seedling		<ul style="list-style-type: none"> Proper drainage to be done and avoid water stagnation
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision against FMD
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide broad-spectrum antihelminthic drugs under vet supervision and recommended doses. Provide adequate clean drinking water+electrolyte and multi-vitamins Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days:

			F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain.
			<ul style="list-style-type: none"> • Remove wet litter
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision • Separate sick animals • The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves • Long hair near the udder/stomach/back legs should be teamed short.
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. Pregnant animals above 7 months of pregnancy should be avoided from vaccination.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा

GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Changlang

No: 43/2017/Tue

Dated: 30th May, 2017



Changlang (HQ): Latitude: 27.1305856/Longitude: 95.7342373/Altitude: 580 masl

Weather summary of the previous week	Weather forecast valid up to 4 th June, 2017
Rainfall = Changlang: 6.0 mm Miao: 17.4 mm	Mostly cloudy weather forecasted with moderate/rather heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 23-29°C and 21-22°C respectively. Expected morning and evening relative humidity will be 95-99% and 70-94% respectively. Wind direction is mainly E with speed range of 2-6 kmph.
Warning	Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting	Citrus scab/ leaf miner	<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> Maintain proper drainage and avoid water stagnation.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth Provide sugar syrup
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-vitamins

			<ul style="list-style-type: none"> • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)




AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Dibang Valley

No: 43/2017/Tue

Dated: 30th May, 2017



Anini: Latitude: 28.2179994/Longitude: 94.7277528/Altitude: 1968 masl

Weather summary of the previous week	Weather forecast valid up to 4 th June, 2017
Rainfall = 36.0 mm (Anini)	Variable with possibility of light to rather heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 14-22°C and 11-13°C respectively. Expected morning and evening relative humidity will be 91-94% and 61-85% respectively. Wind direction is mainly E and ESE with speed range of 3-6 kmph.
Warning:	<ul style="list-style-type: none">Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General recommendations

- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated
- Go for spray operation only on dry/clear weather.
- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> Avoid water stagnation
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Apple	Vegetative	Wooly aphids	<ul style="list-style-type: none"> Swab the stem with mineral oil against wooly aphid.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> In case of rain provide sugar syrup
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-vitamins Vaccination as per the schedule with proper consultation with vet.

			<ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR. BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



भा.क.अनु.प. - पू.प.सं.
ICAR - RCRNEH

AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: East Kameng

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
Rainfall = 10.2 mm	Partly cloudy with possibility of light/moderate rainfall and heavy rainfall on 31 st May. Daily maximum and minimum temperature for the following period is predicted to be 18-22°C and 14-16°C respectively. Expected morning and evening relative humidity will be 95-99% and 70-94% respectively. Expected wind direction is mainly ENE & SE with speed range of 2-4 kmph.
Warning:	<ul style="list-style-type: none"> Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing
- Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting	Citrus scab/ leaf miner	<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> Avoid water stagnation
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Agroforestry			
Toko palm	Nursery		Filling of nursery polybags for raising of takopalm seedlings
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-

			<p>vitamins</p> <ul style="list-style-type: none"> • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Fishery			<ul style="list-style-type: none"> • Best time to prepare the culture pond like cleaning aquatic weeds
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: East Siang

No: 43/2017/Tue

Dated: 30th May, 2017



Pasighat: Latitude: 28.0700/Longitude: 95.3300/Altitude: 153 masl

Weather summary of the previous week	Weather forecast valid up to 4 th June 2017
Rainfall: 17.3 mm; T. Max: 35.0°C; T. Min: 21.0°C; RH: 75-96%	Mostly cloudy with possibility of light/moderate rainfall with heavy rainfall on 31st May. Daily maximum and minimum temperature for the following period is predicted to be 24-29°C and 21-23°C respectively. Expected morning and evening relative humidity will be 92-97% and 82-93% respectively. Expected wind direction is mainly NNE and E with speed range of 2-6 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- Go for low cost polyhouses for year round cultivation of vegetables
- Obtain seeds from reputed/certified/reliable source
- Use fertilizers/pesticides as per recommended doses for the location
- Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides
- ICAR Basar recommend poultry variety VANARAJA for backyard rearing

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting	Citrus scab/ leaf miner	<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> Maintain proper drainage and avoid water stagnation
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> On prolong rain, provide sugar syrup
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Provide adequate clean drinking water+electrolyte and multi-vitamins Give broad-spectrum antihelmintic drugs as per prescribed

			<p>doses.</p> <ul style="list-style-type: none"> Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision
Mithun		FMD	<ul style="list-style-type: none"> Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. Pregnant animals above 7 months of pregnancy should be avoided.
<p>Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247</p>			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Kurung Kumey

No: 43/2017/Tue

Dated: 30th May, 2017



Koloriang: Latitude: 27.9167°/Longitude: 93.3500°/Altitude: 1040 masl

Weather summary of the previous week	Weather forecast valid up to 4 th June 2016
Past weather summary not available	Variable weather with possibility of light to rather heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 20-23°C and 15-16°C respectively. Expected morning and evening relative humidity will be 91-93% and 77-86% respectively. Wind direction is variable with speed range of 2-5 kmph.
Warning	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spray operations should be done only on dry/clear weather or in extreme case use stickers**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**
- **Disinfecting the animal/bird sheds with lime powder/ bleaching powder/ 4%NaOH at regular interval is advocated**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
Pea	Seedling		<ul style="list-style-type: none"> Provide proper drainage to already sown crops
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi 1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting	Citrus scab/ leaf miner	<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> Avoid water stagnation
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
			<ul style="list-style-type: none"> Box must be covered with straw/dry leaves to provide warmth
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Cattle	All stages	FMD	<ul style="list-style-type: none"> 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision Separate sick animals The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves Long hair near the udder/stomach/back legs should be teamed short.
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Provide adequate bedding material (paddy straw, rice husk) to newly born piglets. Animals should be given lukewarm feed and water. All animals should be tethered in a covered shelter. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision

Poultry	All stages		<ul style="list-style-type: none"> • Follow 'All in All out' system of rearing and do not rear birds of different age groups together. • Avoid overcrowding. • Provide adequate clean drinking water+electrolyte and multi-vitamins • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Mithun		FMD	<ul style="list-style-type: none"> • Polyvalent FMD vaccine with serotype O, A and Asia at 6 months of age and repeat after that 6 months interval by sub-cutaneous route. • Pregnant animals above 7 months of pregnancy should be avoided.
Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247			

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**

कृषि मौसम सलाहकार सेवा पत्रिका

Agro-met Advisory Bulletin

(Period: 31st May to 2nd June 2017)

ग्रामीण कृषि मौसम सेवा



GRAMIN KRISHI MAUSAM SEWA
(Integrated Agro-Meteorological Advisory Services)



AMFU, ICAR Research Complex for NEH Region
Arunachal Pradesh Centre, Basar-791101

District: Lohit

No: 43/2017/Tue

Dated: 30th May, 2017



Weather summary of the previous week	Weather forecast valid up to 4 th June, 2017
Rainfall = 21.4 mm (Tezu)	Partly/mostly cloudy weather with possibility of moderate to heavy rainfall. Daily maximum and minimum temperature for the following period is predicted to be 20-23°C and 15-16°C respectively. Expected morning and evening relative humidity will be 81-87% and 47-64% respectively. Wind direction is E with speed range of 2-6 kmph.
Warning:	• Very heavy rainfall is likely over isolated places on 31st May and 1st June

Weather Based Agro-meteorological Advisories Bulletin

General Recommendations

- **Spraying operation should be done only on dry/clear weather**
- **Go for low cost polyhouses for year round cultivation of vegetables**
- **Obtain seeds from reputed/certified/reliable source**
- **Use fertilizers/pesticides as per recommended doses for the location**
- **Preference must be given on use of organic manures/bio-fertilizers & bio-pesticides**
- **ICAR Basar recommend poultry variety VANARAJA for backyard rearing**

Crop	Stage	Pest/Disease	Precautionary Practices to be followed
FIELD CROPS			
Rice	Nursery/transplanting		<ul style="list-style-type: none"> Maintain proper drainage for nursery plot of rice. Seedling in lowland rice nursery may be delayed until further forecast.
Soybean	Sowing		<ul style="list-style-type: none"> Avoid sowing If already sown, maintain proper drainage
Jhum field	Tillering		<ul style="list-style-type: none"> Manual weeding advocated Top dressing of urea one third of recommended dose can be carried out in upland/jhum rice
Maize	Knee height		<ul style="list-style-type: none"> Carry out thinning and earthing up in maize without delay
Finger millet	Sowing		<ul style="list-style-type: none"> Sowing of short duration finger millet varieties like VL-324 & VR-708 can be carried out without delay.
VEGETABLES			
Pre-summer vegetables	Seedling/vegetative		<ul style="list-style-type: none"> Provision of drainage to remove excess water followed by earthing up in vegetables. In cucurbitaceous crops like bottle gourd, cucumber, ridge gourd etc, earthing up to be done followed by manuring; and stacking should be followed and trained on a bower system made of bamboo + sticks. Seed treatment: Before sowing treat seed with trichoderma viridi1% WP @ 10g/kg seed.
HORTICULTURAL CROPS			
Citrus	Fruiting		<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour.
Pineapple	Vegetative		<ul style="list-style-type: none"> Earthing-up and removal of excess suckers for better plant establishment and anchorage.
Ginger/Turmeric	Sowing/germination		<ul style="list-style-type: none"> Avoid water stagnation
Banana	Vegetative		<ul style="list-style-type: none"> Planting of banana making pit maintaining at a spacing of 2.5 X 2.5 meter plant to plant and row to row distance. Denavelling (removal of male bud) to improve bunch weight Bagging of bunches to protect against unusual rainfall.
Orchards			<ul style="list-style-type: none"> Prepare half moon terraces in fruit crops (orange/banana/apple/peach) grown in steep slopes to prevent nutrient loss during heavy downpour. Follow proper drainage and earthing up practices in low land/plain areas to avoid water stagnation in fruit orchards
Large Cardamom	Sowing		<ul style="list-style-type: none"> Should be sown in Primary nursery bed at 2cm deep and 10 cm apart followed by mulching & regular light irrigation. The duration of primary nursery being 3-4 months.
Honey Bees			
Honey bees			<ul style="list-style-type: none"> Provide sugar syrup near the box Add insulation and ventilation systems
LIVESTOCKS			
Livestock	Stage	Disease	Precautionary Practices to be followed
Fishery	Fingerlings		<ul style="list-style-type: none"> Broadcasting of Mustard Oil Cake (MOC) @ 5kg/pond twice daily advocated for healthy fingerlings
Pigs	All stages	FMD (high fever and appearance of mucous membrane of the mouth and foot.)	<ul style="list-style-type: none"> Animals should be given lukewarm feed and water. Keep the houses dry and clean. 1st injection at 6 months of age and 2nd injection at 12 months of age followed by annual vaccination under vet supervision
Poultry	All stages		<ul style="list-style-type: none"> Follow 'All in All out' system of rearing and do not rear birds of different age groups together. Avoid overcrowding. Provide adequate clean drinking water+electrolyte and multi-

			<p>vitamins</p> <ul style="list-style-type: none"> • Give broad-spectrum antihelmintic drugs as per prescribed doses. • Vaccination as per the schedule with proper consultation with vet. <ul style="list-style-type: none"> ➤ Day old chick: HVT Marek disease vaccine, 4-7 days: F/Lasota, 14-18 days: Intermediate plus/IBD vaccine, 35 days: F/Lasota, 6-7 weeks: Chicken embryo adopted fowl pox vaccine and 56-70 days: RD R-2B strain. • Remove wet litter/provide artificial heating and light
Cattle	All stages	FMD	<ul style="list-style-type: none"> • 1st injection at 6-8 weeks of age, 2nd injection after 6 months of 1st injection followed by annual vaccination under vet supervision • Separate sick animals • The animal should be washed with lukewarm water added with little potash (KMnO₄) or neem leaves • Long hair near the udder/stomach/back legs should be teamed short.

Contact Scientists (ICAR) or State Agril Staff or KVK staff for measures against adverse conditions or call at 03795-226237/226247

Issued jointly by ICAR-RC for NEH Region, Arunachal Pradesh Centre, Basar and India Meteorological Department
Contact: - IAAS.BASAR@gmail.com, Phone Number: - 03795-226247/37, -96 (Fax)

**Agrometeorological Field Unit
ICAR, BASAR**