



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Chandel**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Chandel (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 22.0 – 24.0 °C & minimum temp. 7.0 – 9.0 °C. Relative humidity may reach upto 99 % (maximum RH) & 54 % (minimum RH). The average wind speed may be 2-4 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Churachandpur**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Churachandpur (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 25.0 °C & minimum temp. 6.0 – 8.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 48 % (minimum RH). The average wind speed may be 2-4 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Imphal-East**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Imphal-East (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 24.0 °C & minimum temp. 5.0 – 6.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 47 % (minimum RH). The average wind speed may be 0-2 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Imphal-West**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Imphal-West (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 22.0 – 24.0 °C & minimum temp. 5.0 – 7.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 46 % (minimum RH). The average wind speed may be 2-4 km/ hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Senapati**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Senapati (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 21.0 – 23.0 °C & minimum temp. 3.0 – 5.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 47 % (minimum RH). The average wind speed may be 2-3 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to vegetative		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

			10 days till the seedlings get established. - Give light irrigation if the soil is dry.
<b>Animal Sciences</b>			
* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.
<b>Fishery:</b>			
Type of Farming	Stage & Management	Agro-metrological advisories	
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	<ul style="list-style-type: none"> <li>- Fish harvesting can be done.</li> <li>- Stock fish for next year and for this practices mentioned below may be done in small scale.</li> <li>- Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m.</li> <li>- Feeds should be provided 1-2 times a day @ 2-3 % body weight per day.</li> <li>- Well balanced feed fortified with vitamins and minerals should be provided.</li> <li>- Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.</li> </ul>	

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Tamenglong**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Tamenglong (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 22.0 – 24.0 °C & minimum temp. 4.0 – 6.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 46 % (minimum RH). The average wind speed may be 2-4 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Thoubal**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Thoubal (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 24.0 °C & minimum temp. 6.0 – 8.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 51 % (minimum RH). The average wind speed may be 0-4 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Ukhrul**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Ukhrul (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 21.0 – 23.0 °C & minimum temp. 3.0 – 5.0 °C. Relative humidity may reach upto 99 % (maximum RH) & 52 % (minimum RH). The average wind speed may be 2-4 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to vegetative		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

			10 days till the seedlings get established. - Give light irrigation if the soil is dry.
<b>Animal Sciences</b>			
* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.			
Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.
<b>Fishery:</b>			
Type of Farming	Stage & Management	Agro-metrological advisories	
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	<ul style="list-style-type: none"> <li>- Fish harvesting can be done.</li> <li>- Stock fish for next year and for this practices mentioned below may be done in small scale.</li> <li>- Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m.</li> <li>- Feeds should be provided 1-2 times a day @ 2-3 % body weight per day.</li> <li>- Well balanced feed fortified with vitamins and minerals should be provided.</li> <li>- Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.</li> </ul>	

(I. Meghachandra Singh)  
Scientist i/c, GKMS



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

**MANIPUR BULLETIN**

**5<sup>th</sup> December, 2014**

**Bishnupur**

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 6<sup>th</sup> to 10<sup>th</sup> December, 2014 as below:

**Significant past weather for the preceding bi-weekly of Imphal - West (2<sup>nd</sup> to 4<sup>th</sup> December, 2014)**

During past three days sky was mainly clear. **No** rainfall was recorded during past three days. Maximum temperature range observed was 23.5 – 24.3 °C, range of minimum temperature observed was 5.9 – 7.6 °C, mean relative humidity observed in morning and afternoon were 89.3 % and 49.7 % respectively. The average wind speed range was 0.9 – 1.2 km/hr.

**Weather forecast for next five days of Bishnupur (6<sup>th</sup> to 10<sup>th</sup> December, 2014)**

There may be no rain upto Wednesday. Sky will be mainly clear to partly cloudy with maximum temp. of 23.0 – 25.0 °C & minimum temp. 5.0 – 7.0 °C. Relative humidity may reach upto 100 % (maximum RH) & 49 % (minimum RH). The average wind speed may be 2-4 km/hr during next five days.

Name of the crop	Stage	Pest and disease infestation	Agro-meteorological advisories
<b>Field Crops</b>			
Kharif Rice	Post harvesting & storage		<ul style="list-style-type: none"> <li>- For seed, select healthy mature seeds and store in RC-seed bins after proper drying.</li> <li>- To prevent the infestation of stored grains by moths and insects, godowns/ granaries/ bins must be well cleaned and walls be sprayed with 1% Malathion.</li> </ul>
Rapeseed/ mustard	Vegetative		<ul style="list-style-type: none"> <li>- Thinning may be done where the plant population is high.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Field pea	Vegetative		<ul style="list-style-type: none"> <li>- Avoid nipping of young shoots for vegetable.</li> <li>- Give light irrigation if the soil is dry.</li> </ul>
Potato	Vegetative		<ul style="list-style-type: none"> <li>- Give light irrigation if the soil is dry.</li> </ul>
Lentil	Sowing		<ul style="list-style-type: none"> <li>- Lentils may be sown with minimum tillage or zero tillage in rows.</li> </ul>
Rabi maize	Land preparation		<ul style="list-style-type: none"> <li>- Land preparation for sowing rabi maize may be started.</li> <li>- Rabi maize may be sown as intercrop with potato or pulses. Ridges along east-west direction may be prepared after ploughing. Southern face may be sown with maize and northern with pea seeds.</li> </ul>
Storage			<ul style="list-style-type: none"> <li>- Clean and store the seeds in RC-seed bins after proper drying of seeds and charcoal. Check for germination of seeds or fungal growth. After proper drying, store grains/ seeds in well ventilated godowns.</li> </ul>
<b>Horticulture Crops</b>			
Garden Pea	Land preparation to sowing		<ul style="list-style-type: none"> <li>- For mid season pea, thorough ploughing of land but not too fine tilt which will harness soil aeration is required to avoid compactness and also to keep them in aerobic condition of the soil when the rain comes.</li> <li>- As a precautionary measure against wilt (non-filling up of pods or drying up of leaves in lower parts), treat seeds with <i>Trichoderma viride</i> and apply 5-10 kg in soil at the time of sowing and another after 20-30 days.</li> </ul>
Vegetables/ fruits	Soil application		<ul style="list-style-type: none"> <li>- For applying organic formulation in the soil, prepare Sanjivak by mixing 100kg cowdung +100 lit of cow urine + 500g of jiggery + 250 lit of water in a 500 lit capacity closed drum. Ferment for 10 days. Dilute it with 20 times water and sprinkle in one acre of land, either as soil application or along with irrigation water.</li> <li>- For kitchen garden, take 2kg cowdung and 200g oil cake in 20litres capacity plastic bucket and keep it to ferment for 10 days. Dilute it with 20 times water and apply in soil for vegetables and flowers.</li> <li>- To avoid unpleasant smell, it may be kept in open for 1-2 days.</li> </ul>
Onion	Transplanting to		<ul style="list-style-type: none"> <li>- Maintain 50-60 days old seedling transplanted. Transplant at a spacing</li> </ul>



**ICAR Research Complex for N.E.H. Region  
Manipur Centre, Lamphelpat, Imphal - 795 004**



**“Integrated Agromet Advisory Services”**

	vegetative		of 10-20cm or 10-15cm as per choice. Moist the bed before transplanting. Cover the seedlings only upto the root portion. Slight watering during and evening is very much necessary for at least 7-10 days till the seedlings get established. - Give light irrigation if the soil is dry.
--	------------	--	--

**Animal Sciences**

\* Cover side walls of birds/ animals sheds with polythene sheet or hessian cloth to protect from cold weather.

Name of the animal	Growth stage	Disease/ infestation	Agro-meteorological advisories
<b>Poultry</b>	All susceptible age groups (as and when required)	Coccidiosis	Give preventive dose of Coximar @ 1g/ lit of drinking water for 7 days along with glucose.
		Management	- Mix deactivated lime powder @ 2 kg per 100 sq. ft. area in litter material. - Increase energy feed upto 5% (like maize, rice bran, etc.). - Add antibiotic, Hostacycline/ Duaprin @ 2 g/ lit of drinking water for 7 days along with vitamin B & vitamin A.
<b>Sheep/ Goat</b>	Above 3 months of age (as and when required)	Tetanus	- In pregnant animals, inject Tetanus Toxoid @ 0.5ml at 3.5 months of pregnancy.
		Ectoparasite	Use Butox liq. @ 2ml/ 4 litres of water and wash the animal with the solution.
		Deworming	- Give Albomac suspension @ 5ml per adult.
<b>Pig</b>	Above 3 months of age (as and when required)	Skin infection	Give Ivermectin (Indectin tab.) @ 10mg/33kg body weight.
		Deworming	- Deworm using Piperazine citrate @ 30 ml per adult animal once.
		Ticks	Add Butox liquid @ 2 ml/ lit of water and use the solution twice a week while brushing the animal.
<b>Cattle / Buffalo/ Mithun</b>	Above 3 months of age (as and when required)	B.Q.	- Vaccinate against B.Q. @ 1ml orally/ sub- cutaneously especially in young stock.
		F.M.D.	- Vaccinate against F.M.D.
		Ticks/ Wounds	Topicure spray may be used twice daily on the affected parts.

**Fishery:**

Type of Farming	Stage & Management	Agro-metrological advisories
<b>Carp/ Catfish culture</b>	Harvesting and stock maintenance	- Fish harvesting can be done. - Stock fish for next year and for this practices mentioned below may be done in small scale. - Ponds should be free from unwanted vegetation and insects and water depth should be maintained at 1.0 - 1.5m. - Feeds should be provided 1-2 times a day @ 2-3 % body weight per day. - Well balanced feed fortified with vitamins and minerals should be provided. - Apply Green AQ @ 4litre/ ha or lime @ 200kg/ ha. Maintain good quality of pond water.

(I. Meghachandra Singh)  
Scientist i/c, GKMS