



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Bishnupur

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Bishnupur (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Sunday with mainly cloudy to overcast sky and maximum temp. 28 - 32 °C & minimum temp. 23 - 25 °C. Relative humidity may reach up to 99 % (max RH) & 98 % (min RH). The average wind speed may be 2 - 4 km/hr.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

(I. Meghachandra Singh)
Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Chandel

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Chandel (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Sunday with mainly cloudy to overcast sky and maximum temp. 28 - 33 °C & minimum temp. 23 - 25 °C. Relative humidity may reach up to 99 % (max RH) & 98 % (min RH). The average wind speed may be 2 - 4 km/hr.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

**(I. Meghachandra Singh)
Scientist i/c, GKMS**



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Churachandpur

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Churachandpur (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Sunday with mainly cloudy to overcast sky and maximum temp. 28 - 32 °C & minimum temp. 24 - 25 °C. Relative humidity may reach up to 99 % (max RH) & 98 % (min RH). The average wind speed may be 0 - 4 km/hr.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @ 3ml/15 lit of water or Lambdacythrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

(I. Meghachandra Singh)
Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Senapati

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Senapati (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Wednesday with mainly cloudy to overcast sky and maximum temp. 27 - 32 °C & minimum temp. 21 - 24 °C. Relative humidity may reach up to 99 % (max RH) & 96 % (min RH). The average wind speed may be 2 - 3 km/hr.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythothrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

(I. Meghachandra Singh)
Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Tamenglong

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Tamenglong (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Wednesday with mainly cloudy to overcast sky and maximum temp. 27 - 32 °C & minimum temp. 23 - 24 °C. Relative humidity may reach up to 99 % (max RH) & 98 % (min RH). The average wind speed may be 0 - 3 km/hr.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythothrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

(I. Meghachandra Singh)
Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Thoubal

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Thoubal (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Sunday with mainly cloudy to overcast sky and maximum temp. 28 - 33 °C & minimum temp. 23 - 25 °C. Relative humidity may reach up to 99 % (max RH) & 98 % (min RH). The average wind speed may be 0 - 4 km/hr.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

(I. Meghachandra Singh)
Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Ukhrul

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Ukhrul (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Wednesday with mainly cloudy to overcast sky and maximum temp. 27 - 33° C & minimum temp. 22 - 24 °C. Relative humidity may reach up to 99 % (max RH) & 97 % (min RH). The average wind speed may be 2 - 5 km/hr.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

**(I. Meghachandra Singh)
Scientist i/c, GKMS**



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Imphal East

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Imphal East (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Wednesday with mainly cloudy to overcast sky and maximum temp. 27 - 33 °C & minimum temp. 22 - 24 °C. Relative humidity may reach up to 99 % (max RH) & 96 % (min RH). The average wind speed may be 0 - 3 km/hr.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythothrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

(I. Meghachandra Singh)
Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****11th August, 2017**Imphal West

IAAS, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and Northern Hemisphere Analysis Centre, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 12th to 16th August, 2017 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (8th to 10th August, 2017) Observed to mainly cloudy to overcast sky and 4.9 mm rainfall was recorded during past three days. Maximum temperature range of 27.3 to 28.2 °C and minimum temperature range of 21.6 to 22.8 °C with mean relative humidity 94.3 % max and 82.0 % min was recorded in the past three days.</p>	<p>Weather forecast for next five days of Imphal West (12th to 16th August, 2017) Likely to have heavy to very heavy rains at isolated places upto Wednesday with mainly cloudy to overcast sky and maximum temp. 27 - 32 °C & minimum temp. 23 - 24 °C. Relative humidity may reach up to 99 % (max RH) & 97 % (min RH). The average wind speed may be 0 - 3 km/hr.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

State	Weather Forecast/heavy rainfall warning issued on 11.08.17 (morning)	Advisory
Nagaland, Manipur, Mizoram & Tripura.	Rainfall likely at most/many places during next 5 days along with heavy to very heavy rain at isolated places on 11 th August.	Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Arrange for adequate drainage to avoid water stagnation in the crop fields

Field Crops

* Crops must be exposed to sun for at least three hours after spraying of the following insecticide and pesticides. Hence they must be applied if there is probability of no rain for at least three hours.

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Tillering (kharif rice)		- Off-type plants must be removed. - After rains, drain out water and apply 33kg of urea as top dress after 30days of transplanting. Drain in water after 48hrs of top-dressing.
		Bacterial leaf blight	- Streptocycline @ 2g/15 litres of water (at least three pumps per sangam) after rains.
		Any sucking insect Case worm, borer, leaf folder, thrips	- Imidachlopid @ 8ml/ 15 litres of water after rains. - Coragen @3ml/15 lit of water or Lambdacythothrin @ 15ml/ 15 litre of water (at least three pumps per sangam) after rains.
Maize	Cob formation		- Weeding may be done after rains to avoid insect pest in the crop field and also avoid water logging in the crop field. - Construct proper drainage to avoid water logging.
Soybean / Groundnut	Vegetative to flowering		- Avoid leaf eating caterpillars/ insects at least upto 60 days after sowing. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
		Tikka disease	- Spray Carbendazim @ 0.5g / lit of water after rains to control Tikka disease in groundnut.
Blackgram & Rice bean	Vegetative to flowering		- Avoid hairy caterpillars during vegetative stage. - Construct proper drainage to avoid water logging.
		hairy caterpillars	- For control of red hairy caterpillars, dust 5% Malathion @ 20 kg/ ha to control it or spray Dichlorovos 7.6 EC 0.05% (0.6ml/ lit of water) after rains.
Seeds storage	Post harvest management		- For seed purpose dry, clean and store the seeds in RC-seed bins after proper drying of seeds along with charcoal at 80:20. Check for moisture content or weight of charcoal and dry if required. - For grains, store grains in well ventilated godowns after drying proper. Spray Malathion 1% on the walls of stores.

Horticulture Crops

Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Turmeric	Vegetative	Leaf Spot and leaf blotch	- Spray DM-45 @ 2.0 g/ lit of water as prophylactic measure against Leaf spot and Leaf blotch at 7 days interval after rain. - Drain out excess water to avoid water logging to avoid soft rot.
Cauliflower and Tomato	Nursery of mid season variety		- Use temporary rain shelter facility/ low cost polyhouse for raising nursery. Protect nursery from damping off. Fumigate the soil with formaldehyde drench the soil with Captan 50WP.



ICAR Research Complex for N.E.H. Region

Manipur Centre, Lamphelpat, Imphal - 795 004

“Gramin Krishi Mausam Seva”



Cauliflower	Transplanting to early vegetative (early variety)		- Transplant healthy seedlings of early cauliflower (variety like Early Himlata/ White Shot/ White treasure) at a spacing of 45cm x 45cm.
King Chilli	Fruiting	Fruit rot and aphid	- Spray Redomil Gold @ 2g/lit water + Imidacloprid @ 3ml/10lit of water after rains.
Onion	Sowing/ planting		- This is the season for sowing of onion in the upland areas.
Banana	For old orchard		- Remove unwanted suckers. After 30-35 days of harvesting fruits, mother plant must be cut at a height of 0.6m or 2 ft. Maintain only 3 plants per pit.

Animal Sciences

*** Provide clean drinking water for birds and animals.
- Check for water leakage in sheds of birds and animals.**

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Turkey/ Duck)	All age groups	Deworming	- Use Albendazole liquid @ 20-40 ml for 100 adult birds and repeat it next day with half the dose at regular interval of 3 weeks.
			- Bleaching powder may be spread near farm premises. - Avoid fungal infested feed. - Use dry litter material in the shed, if moist apply lime powder @ 1kg per 100 ft ² area.
Sheep/ Goat	All age groups	Vaccination	- Sheep should be vaccinated against B.Q./ H.S.
Pig	All age groups except the young ones	De-worming	- De-worming of animals should be done using Albendazole @7.5 mg/kg b.wt.
		Vaccination	- Vaccinate against FMD.
		Management	- Provide adequate amount of wholesome water (drinking). - Essential body salts may be provided.
Cattle	All age groups excepts the young ones	B.Q. & H.S.	Vaccinate against B.Q., F.M.D. & H.S.
		Deworming	Use Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		Slush & Flood Management	- Provide clean and wholesome water. Keep animals in safe place. Avoid grazing at the periphery of flood. Animals should be protected against illness caused due to damp weather. Mineral mixtures/vitamins should be fed at this time. Do not allow to drink flood water.
		Milk fever in Milch cow	70-100ml of a mixture of Ca+ and Phosphorus daily @ 5-7days.

Fishery:

*** Encircle fish ponds with happa net to avoid escape of fishes during rainy days.**

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Nurseries/ grow-out	Pond and water	Maintain minimum depth of 1.0 - 1.5 m and pH at 7.0 – 7.5. Drain out excess water accumulated in fish pond due to rain.
		Feed	Provide feed 2-3 times @ 5-10% body weight per day. Provide good feed having 30% crude protein fortified with vitamins and mineral.
		Fish	For semi-intensive farming, 8000-10000 fingerlings can be stocked in one hectare pond. - Stocking ratio may be maintained at 40% of surface feeder, 30% column feeder and 30% bottom feeder. - Sampling of fish for growth, disease should be done regularly by netting.
Integrated Fish Farming	Fish	Same as grow out fish management	Same as grow out fish farming management.
	Pig		About 30-40 two months old pigs may be kept for proper fertilization of the one ha pond. Pigs are feed on mash at an average of 1kg/day. Green grasses or animal folder is also provided. Grass with interlocked soil in the root system (sod) is provided once a week to avoid mineral deficiency.
	Duck		200-400 ducks of 2-3 months old are sufficient to adequately fertilize a 1 ha pond.

Apply Agri lime in the fish culture pond after the water level reaches normal depth after flood @ 30-35 kg /ha at 30days interval.

**(I. Meghachandra Singh)
Scientist i/c, GKMS**