



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****1st June, 2018**Bishnupur

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Bishnupur (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 32 °C & minimum temp. 21-22 °C. Relative humidity may reach up to 100 % (max RH) & 67 % (min RH). The average wind speed may be 0-2 km/hr.</p>
---	--

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension @60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given @5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



MANIPUR BULLETIN

1st June, 2018

Chandel

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Chandel (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 28-29 °C & minimum temp. 18 °C. Relative humidity may reach up to 96 % (max RH) & 58 % (min RH). The average wind speed may be 3 km/hr.</p>
---	---

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension @60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given @5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



MANIPUR BULLETIN

1st June, 2018

Churachandpur

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Churachandpur (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 31-32 °C & minimum temp. 19 °C. Relative humidity may reach upto 100 % (max RH) & 70 % (min RH). The average wind speed may be 0-2 km/hr.</p>
---	---

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given@5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



MANIPUR BULLETIN

1st June, 2018

Imphal East

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Imphal East (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 31-32 °C & minimum temp. 19-20 °C. Relative humidity may reach up to 100 % (max RH) & 51 % (min RH). The average wind speed may be 2 km/hr.</p>
---	---

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension @60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given @5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



MANIPUR BULLETIN

1st June, 2018

Imphal West

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Imphal West (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 31-32 °C & minimum temp. 19-20 °C. Relative humidity may reach up to 100 % (max RH) & 63 % (min RH). The average wind speed may be 2 km/hr.</p>
---	---

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery an land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given@5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



MANIPUR BULLETIN

1st June, 2018

Senapati

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Senapati (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 27-28 °C & minimum temp. 16-17 °C. Relative humidity may reach upto 100 % (max RH) & 63 % (min RH). The average wind speed may be 2-3 km/hr.</p>
---	---

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



<i>Animal Sciences</i>			
Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

<i>Fishery</i>			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given@5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****1st June, 2018**Tamenglong

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Tamenglong (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 28-29 °C & minimum temp. 18 °C. Relative humidity may reach upto 100 % (max RH) & 58 % (min RH). The average wind speed may be 2 km/hr.</p>
---	--

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension @60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given @5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****1st June, 2018**Thoubal

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Thoubal (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 32 °C & minimum temp. 21-22 °C. Relative humidity may reach up to 99 % (max RH) & 67 % (min RH). The average wind speed may be 2-4 km/hr.</p>
---	---

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Tomato/ Chilli	Flowering & fruiting to harvesting		- Retain mature and healthy fruits of chilli for seeds.
		Late Blight	- Harvest mature ones and spray Dithane M-45 @ 2g/lit of water.

Animal Sciences

Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension @60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery

Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given @5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”

**MANIPUR BULLETIN****1st June, 2018**Ukhrul

GKMS at AMFU- Lamphelpat, ICAR-RC Manipur Centre in association with India Meteorological Department, Regional Meteorological Centre, Guwahati and India Meteorological Department, New Delhi, Ministry of Earth Science, Govt. of India reports the weather forecasts for the next five days, i.e., from 2nd to 6th June, 2018 as below:

<p>Significant past weather for the preceding bi-weekly of Imphal - West (29th to 31st May, 2018) Observed 4.5 mm rainfall with partly cloudy to overcast sky during past three days. Maximum temperature recorded ranged from 30.7 to 31.5 °C and minimum of 20.4 to 20.9 °C with mean relative humidity 87.7 % max and 60.0 % min was recorded in the past three days with average wind speed was 1.9- 4.9 km/ hr.</p>	<p>Weather forecast for next five days of Ukhrul (2nd to 6th June, 2018) Likely to have heavy rains at isolated places during next five days and partly cloudy to overcast sky with maximum temp. 28-29 °C & minimum temp. 16-17 °C. Relative humidity may reach upto 98 % (max RH) & 56 % (min RH). The average wind speed may be 2-3 km/hr.</p>
---	--

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Panicle initiation to flowering (Pre-kharif rice)		- Maintain water level at 5-8 cm. - Monitor rice fields regularly for insect pest infestation.
	Nursery and land preparation of main field (Kharif rice)		- Raise kharif rice nursery. Varieties: RC Maniphou-7, RC Maniphou- 10 and RC Maniphou- 13. - Apply well decomposed FYM/ compost @ 1t/ 700 sq. m. at final leveling of nursery bed. After 7-10 days of sowing, apply 17-20 kg Urea and 3.5 kg of MOP per 700 m ² along with 2 kg Carbofuran 3G, 2g Streptocycline and 20g Carbendazim. Maintain water level just to submerge green sprouts. - First plough of the rice field may be done. Submerge the soil for 15-20 days.
Kharif Maize	Sowing		- Sowing may be done.
Soybean	Land preparation to Sowing		- Sow seed after rains @ 40-45 kg/ha or 10-11.3 kg of seed/ sangam and the recommended variety is J.S.335. Sowing may be done at spacing of 45cm x 30cm. Treat seeds with Nativo@2g/kg seeds or with Bavistin@2.5g/kg seed or with Thiram.
Groundnut	Land preparation to Sowing		- Seed rate is 100 – 110 kg/ha and the recommended varieties are ICGS-75 & JL-24. Sow seeds at spacing 30cm x 15cm after rains. Treat seeds with Imidacloprid @ 2ml/ litre water or Bavistin @ 2.5g/ kg seed or with Thiram.
Blackgram/ Rice Bean	Land preparation to Sowing		- Prepare land after rains. Sowing may be done by broadcasting or dibbling. Plot must be kept weed free to avoid insect/ pest.
Seeds storage	Post harvest management		For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. For grains, store grains in well ventilated godowns after drying properly. Spray Malathion 1% on the walls of stores.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Brinjal	Flowering to harvesting	Shoot & fruit borer	- Prune dropping shoots and burn it. Destroy the fruit along with borer. - Spray Cypermethrin @ 2ml/ lit of water.
Cucurbits (Water-melon, pumpkin, sponge gourd, etc.)	Flowering & fruiting to harvesting	Powdery mildew or Downey mildew	- Harvest mature fruits and spray Bayleton (Triademeton 25WP) @ 0.7 g/ lit of water may be done twice at 10 & 14 days intervals respectively against Powdery mildew or Downey mildew especially in damped or moist area. - Drain out accumulated/ excess water in the field. - Sow cucurbits in upland area or under polyhouse.
Ginger/ Turmeric	Vegetative		Crops that were planted in the month of March may be top dressed with 25kg DAP + 10 Kg MOP per sangam in upland areas.
Colocassia	Vegetative		- 45-50 days old plants may be top dressed with 25 kg DAP/ sangam. - Prophylactic measures against leaf blight should be taken up. Spray Redomil@ 1.5g/litre of water or Mancozeb @ 2gm/lit of water twice or thrice at 10 days interval after this wet spell.



ICAR Research Complex for N.E.H. Region
Manipur Centre, Lamphelpat, Imphal - 795 004
“Gramin Krishi Mausam Seva”



Animal Sciences			
Name of bird/ animal	Stage	Disease and Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	• Ranikhet disease	• Strongly advised to vaccinate against Ranikhet disease. Vaccinate the flock using Lasota strain vaccine through ocular route if the age of chick is less than 21 days and through nasal route if it is older than 21 days.
		• Endoparasites	• De-worm using Levamisole Hydrochloride. Potash granules may be added in the drinking water of poultry.
		• Coccidiosis	• Give Amprolium @ 2 gms/ litre of drinking water for 5-7 days.
Sheep/ Goat	All age groups	• Vaccination	• If the animals have not been vaccinated against F.M.D., B.Q., H.S., Enterotoxaemia, etc. do it in this month.
		• Management	• Fly repellent must be used in the premises of the sheds. • Protect from thunder showers. Protect from after winds (loo).
Pig	All age groups except the young ones	• De-worming	• De-worm with Albendazole @7.5 mg/kg b.wt.
		• Vaccination	• Vaccinate against FMD and Swine Fever.
		• Skin disease	• Sprayer of <i>Anti-Psoroptic</i> may be used.
		• Management	• Breeding must be advocated continuously. • Protect from thunder showers especially during night.
Cattle	All age groups excepts the young ones	• Vaccination	• Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated. • Cases of H.S. may be encountered during this period • Immediate treatment using appropriate drugs with the help of Consultant must be advocated to safe the animals.
		• Deworming	• De-worm with Albomac suspension@60-90ml for 300kg body weight or 20-30ml for 100 kg body weight.
		• Care & Management	• Mixture of mineral/ salts should be added in the feed/water to avoid loss of essential body salts. Avoid the cattle from chiting/ biting by flies. Adopt tick control measures

• Fodder cultivation: Berseem/ oat crop may be sown immediately.

Fishery			
Type of Farming	Stage	Management	Agro-metrological advisories
Carp culture	Brooders	Pond and water	Ensure good quality/ clean water free from unwanted plants and animals with a minimum depth of 1.5 m. Fish feed pellet or rice bran & oil cakes (in ratio 1:1) should be provided @ 3-5% body weight per day in 2-3 splits. Spray CIFAX on the surface of the pond water. Male and female brooders must be kept in separate tanks.
	Breeding	pond	Fish hormones like ovaprim, ovatide or Wova-FH can be injected in both the brooders @ 0.5 - 0.7 ml/kg of fish.
	Hatcheries	water	Hatching can be done in traditional hapas having outer & inner hapas or circular hatcheries. A mild current with fine shower is needed for circular hatcheries. Feed should not be given for 3 days.
	Nurseries	ponds	- Stock 20-30 lakhs spawns in one hectare water. Water depth should be more than 1.0 m. Feeding should be given@5-10% body weight/day.

(I. Meghachandra Singh)
 Scientist i/c, GKMS