

**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018

District: Bishnupur

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Bishnupur (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 26 °C & minimum temp. 15-16 °C. Relative humidity may reach upto 97 % (max) and 61 % (min). The average wind speed may be 2 km/hr.
<b>General Advisories:</b>	
* Harvested crop must be kept under rain shelter sheds until sun shines.	
* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.	
* Cover the nursery bed to avoid birds or insect till germination.	
* Mulching may be done wherever possible to retain soil moisture for longer period.	

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety – M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with Rhizobium leguminosarum or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with T. harzianum. Check that the new sprouts/ young plants in nursery beds are not damaged.
Animal Sciences			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.
Fisheries			
Carp culture			- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
Nodal Officer, GKMS

**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018

District: Chandel

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Chandel (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 26 °C & minimum temp. 15-16 °C. Relative humidity may reach upto 94 % (max) and 48 % (min). The average wind speed may be 3 km/hr.
<b>General Advisories:</b>	
* Harvested crop must be kept under rain shelter sheds until sun shines.	
* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.	
* Cover the nursery bed to avoid birds or insect till germination.	
* Mulching may be done wherever possible to retain soil moisture for longer period.	

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety - M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with Rhizobium leguminosarum or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with T. harzianum. Check that the new sprouts/ young plants in nursery beds are not damaged.
Animal Sciences			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.
Fisheries			
Carp culture			- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
Nodal Officer, GKMS

**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018

District: Churachandpur

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Churachandpur (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 26 °C & minimum temp. 15-16 °C. Relative humidity may reach upto 98 % (max) and 45 % (min). The average wind speed may be 3-4 km/hr.
<b>General Advisories:</b> * Harvested crop must be kept under rain shelter sheds until sun shines. * Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent. * Cover the nursery bed to avoid birds or insect till germination. * Mulching may be done wherever possible to retain soil moisture for longer period.	

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety - M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.

Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with Rhizobium leguminosarum or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with T. harzianum. Check that the new sprouts/ young plants in nursery beds are not damaged.

Animal Sciences			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.

Fisheries	
Carp culture	- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.



**MANIPUR AGROMET ADVISORY BULLETIN**

Date of issue: 6<sup>th</sup> November, 2018

District: Imphal East

<p><b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3<sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 – 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days &amp; average wind speed was 1.6 – 2.7 km/hr.</p>	<p><b>Weather forecast for next five days of Imphal East (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 26 °C &amp; minimum temp. 15-16 °C. Relative humidity may reach upto 94 % (max) and 48 % (min). The average wind speed may be 3-5 km/hr.</p>
<p><b>General Advisories:</b></p> <ul style="list-style-type: none"> <li>* Harvested crop must be kept under rain shelter sheds until sun shines.</li> <li>* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.</li> <li>* Cover the nursery bed to avoid birds or insect till germination.</li> <li>* Mulching may be done wherever possible to retain soil moisture for longer period.</li> </ul>	

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety – M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in <i>RC-Seed Bin</i> after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.

Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with <i>Rhizobium leguminosarum</i> or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with <i>T. harzianum</i> . Check that the new sprouts/ young plants in nursery beds are not damaged.

Animal Sciences			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.

Fisheries	
Carp culture	<ul style="list-style-type: none"> <li>- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming.</li> <li>- Harvesting of fishes may be done.</li> <li>- Select disease free fishes to be stocked as brooder.</li> </ul>

(I. Meghachandra Singh)  
Nodal Officer, GKMS



## MANIPUR AGROMET ADVISORY BULLETIN

Date of issue: 6<sup>th</sup> November, 2018

District: Imphal West

<p><b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3<sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days &amp; average wind speed was 1.6 - 2.7 km/hr.</p>	<p><b>Weather forecast for next five days of Imphal West (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 26 °C &amp; minimum temp. 15-16 °C. Relative humidity may reach upto 94 % (max) and 48 % (min). The average wind speed may be 4 km/hr.</p>
<p><b>General Advisories:</b></p> <ul style="list-style-type: none"> <li>* Harvested crop must be kept under rain shelter sheds until sun shines.</li> <li>* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.</li> <li>* Cover the nursery bed to avoid birds or insect till germination.</li> <li>* Mulching may be done wherever possible to retain soil moisture for longer period.</li> </ul>	

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety – M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.

Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with Rhizobium <i>leguminosarum</i> or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with T. <i>harzianum</i> . Check that the new sprouts/ young plants in nursery beds are not damaged.

Animal Sciences			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.

Fisheries	
Carp culture	- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
Nodal Officer, GKMS

**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018District: *Senapati*

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Senapati (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 24-25 °C & minimum temp. 14-15 °C. Relative humidity may reach upto 98 % (max) and 56 % (min). The average wind speed may be 3-4 km/hr.
<b>General Advisories:</b> <ul style="list-style-type: none"> <li>* Harvested crop must be kept under rain shelter sheds until sun shines.</li> <li>* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.</li> <li>* Cover the nursery bed to avoid birds or insect till germination.</li> <li>* Mulching may be done wherever possible to retain soil moisture for longer period.</li> </ul>	

<b>Field Crops</b>			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety - M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in <i>RC-Seed Bin</i> after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.
<b>Horticulture Crops</b>			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with <i>Rhizobium leguminosarum</i> or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
<b>Animal Sciences</b>			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
Cattle	All age groups excepts the young ones	Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
		Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.
<b>Fisheries</b>			
Carp culture			- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
 Nodal Officer, GKMS

**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018

District: Tamenglong

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Tamenglong (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 24-25 °C & minimum temp. 14-15 °C. Relative humidity may reach upto 96 % (max) and 52 % (min). The average wind speed may be 3-4 km/hr.
<b>General Advisories:</b>	
* Harvested crop must be kept under rain shelter sheds until sun shines.	
* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.	
* Cover the nursery bed to avoid birds or insect till germination.	
* Mulching may be done wherever possible to retain soil moisture for longer period.	

Field Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety – M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.
Horticulture Crops			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with Rhizobium <i>leguminosarum</i> or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with T. <i>harzianum</i> . Check that the new sprouts/ young plants in nursery beds are not damaged.
Animal Sciences			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.
Fisheries			
Carp culture			- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
Nodal Officer, GKMS

**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018**District: Thoubal**

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Thoubal (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 26 °C & minimum temp. 15-16 °C. Relative humidity may reach upto 97 % (max) and 53 % (min). The average wind speed may be 0-4 km/hr.
<b>General Advisories:</b>	
* Harvested crop must be kept under rain shelter sheds until sun shines.	
* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.	
* Cover the nursery bed to avoid birds or insect till germination.	
* Mulching may be done wherever possible to retain soil moisture for longer period.	

<b>Field Crops</b>			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety – M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in <i>RC-Seed Bin</i> after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.
<b>Horticulture Crops</b>			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with <i>Rhizobium leguminosarum</i> or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Potato	Planting to vegetative		- Planting of potato may be done at a spacing of 60cm x 15-20cm. Seed rate is 25-30 quintals/ hectare. Treat seed tubers with Sixer (Carbendazim + Mancozeb) @ 2 g/lit of water or Carbaryl 50 WP or Ridomil @ 2 gm/kg seed/ tubers for 20 minutes and let it dry in shed before planting.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
Onion	Nursery raising		- Apply Sixer during final land preparation of nursery plot as preventive measures against soil borne diseases. For 25 sq. m area, we need ½ kg seeds. Treat seeds with <i>T. harzianum</i> . Check that the new sprouts/ young plants in nursery beds are not damaged.
<b>Animal Sciences</b>			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.
<b>Fisheries</b>			
Carp culture			- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
Nodal Officer, GKMS



**MANIPUR AGROMET ADVISORY BULLETIN**Date of issue: 6<sup>th</sup> November, 2018

District: Ukhrul

<b>Significant past weather for the preceding bi-weekly of Imphal West (2<sup>nd</sup> to 5<sup>th</sup> Nov, 2018)</b> Observed 0.4 mm rainfall with mainly clear to partly cloudy sky but mainly cloudy during afternoon hours of 3 <sup>rd</sup> Nov. Maximum temperature recorded ranged from 25.3 - 27.0 °C and minimum of 11.5 - 13.1 °C with mean relative humidity 89.3 % max and 63.5 % min was recorded in the past four days & average wind speed was 1.6 - 2.7 km/hr.	<b>Weather forecast for next five days of Ukhrul (7<sup>th</sup> to 11<sup>th</sup> Nov, 2018)</b> Likely to have no rains upto Sunday and partly cloudy to mainly cloudy sky with maximum temp. 24-25 °C & minimum temp. 14-15 °C. Relative humidity may reach upto 96 % (max) and 52 % (min). The average wind speed may be 3 km/hr.
<b>General Advisories:</b>	
* Harvested crop must be kept under rain shelter sheds until sun shines.	
* Sowing of seeds for rabi season may be done as early as possible so that the available moisture may be utilized to maximum extent.	
* Cover the nursery bed to avoid birds or insect till germination.	
* Mulching may be done wherever possible to retain soil moisture for longer period.	

<b>Field Crops</b>			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Rice	Harvest and post-harvest stage		- Harvest mature crops.
Rapeseed/ mustard	Sowing		- After harvesting rice, zero tillage rape seed may be sown when the field is slightly in muddy condition. Apply 65kg Urea + 188kg SSP + 50kg MOP as basal dose before sowing. Recommended variety - M-27. Seed rate is 10-12 kg/ ha. Treat seeds with Mancozeb @2g/kg seed for the control of white rust and Downey Mildew at seedling stage.
Field pea	Land preparation to sowing		- Land preparation may be done. Sowing may be done at a spacing of 30cm X 10-15cm. Seed rate is 75 kg/ hectare.
Seeds storage	Post-harvest management		- For seed purpose store the seeds in RC-Seed Bin after proper drying and cleaning of seeds along with dry charcoal at 80:20 w/w ratios. Check for moisture content or weight of charcoal and dry if required. - Clean godowns and spray with 1% Malathion on the walls before storage; also ensure that it is well ventilated. Avoid fungal growth and pest rodent in store.
<b>Horticulture Crops</b>			
Name of the crop	Stage	Pest and disease	Agro-meteorological advisories
Garden pea	Sowing/ Land preparation		-Sowing of Garden pea may be done at a spacing of 10cm x 20cm. Before sowing, treat seeds with <i>Rhizobium leguminosarum</i> or follow soil inoculation with phosphor-bacterium. Maintain the depth of sowing to 2-3cm. Avoid shade area while sowing.
Cole crops	Vegetative (Mid-season var.) and nursery (late var.)		- Apply 50 kg urea in two splits dose after 20-40 days of transplanting for mid-season crops in vegetative stage. - Monitor for insect pest infestation in nursery plot.
Ginger/ Turmeric	Harvesting		- 9 months old crops may be harvested. Harvested crop must be kept under rain shelter sheds for curing.
<b>Animal Sciences</b>			
Name of bird/ animal	Stage	Disease/ Management	Agro-meteorological advisories
Poultry (Chicken/ Duck)	All age groups	Coccidiosis	Give Coccimar @ 2g/litre of drinking water for 5-7 days along with glucose.
		Deworming	Use Ventrimizole @ 1g in 1 litre of drinking water for 3 days.
Sheep/ Goat	All age groups	Deworming	- Give Ranizole suspension @ 3ml/ 10 kg body weight.
		Fascioliosis	Give Tolzan-F @ 15ml orally/ kg b. wt. It must be associated with vitamins like Livozyme @ 50ml twice daily.
Pig	All age groups except the young ones	Deworming	Give Morantel citrate tablet (Banminth) @15mg/ kg body weight. Use Butox liquid @ 2 ml/ lit of water twice a week while brushing the animal.
		Vaccination	Vaccinate against B.Q., F.M.D. & H.S. if not yet vaccinated.
Cattle	All age groups excepts the young ones	Tick	Use Butox liquid @ 2ml/ litre water twice a week while brushing animals.
		Deworming	De-worm with Nilzan/ Tolzan-F @20-30ml for 100 kg body weight.
<b>Fisheries</b>			
Carp culture			- Maintain water level at a depth of 1.0m to 1.5m. Ensure good quality water in fish culture ponds by liming. - Harvesting of fishes may be done. - Select disease free fishes to be stocked as brooder.

(I. Meghachandra Singh)  
Nodal Officer, GKMS