



Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Dimapur District

Bulletin No:100/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 25°C to 26°C and 14°C to 15°C, respectively.
- Relative humidity varied from 40% to 88%.
- Wind speed ranged from 3 to 4 kmph

- NDVI for Nagaland

Weather forecast valid upto 27th Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 24°C - 25°C and the **min temp** 11°C to 12°C
- **Sky is likely to be clear** the coming week
- **Relative Humidity** is likely to range from 37% to 86%.
- **Wind speed** may reach upto 3- 4 kmph
Wind direction will be easterly

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties

Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

Citrus			Regular monitoring for trunk borer infestation. Diseased and senile branches should be removed
Vegetables			<i>Intercultural operations in vegetables are advised to remove the weeds.</i>
Cauliflower, cabbage,			<i>Transplanting of cabbage and cauliflower should be done at a spacing – 30 x 60cm for small varieties. 45 x 60 cm for large varieties Keep the field free from weed. Hand hoeing should be done after transplanting</i>
Litchi			<i>Mulch the tree basin, protect young litchi plants from cold and suppress new flush by application of growth inhibitors</i>
Ginger / turmeric	Maturity stage		<i>For Ginger and turmeric, field must be inspected daily for disease appearance. While inspecting, the healthy plants/plots must be marked and kept for planting in the next season. Marking has to be done properly otherwise when plant matures and gets dried up, it will be very difficult to find out the marked plots. Earthing up should be done</i>
Livestock			
Provide clean and cool drinking water to the animals			
Poultry			<ol style="list-style-type: none"> 1. Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials 2. Deworm the birds once every 2 months 3. Sick birds showing respiratory distress, drooping of head or any other illness should be promptly segregated and the entire flock be treated against the illness.
Piggery			<ol style="list-style-type: none"> 1. Protect the animals from cold draft using gunny bags and other locally available materials 2. Provide protein rich ration and ad libitum clean drinking water 3. Pregnant sows can be dewormed only with fenbendazole 4. To protect piglet anemia in newly born piglets provide iron dextran injection on 4th or 14th day after birth or alternately provide ferrous sulphate powder for 14 days
Fisheries			



If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha.
If water depth is less than 1m harvest 50% of the stocked fish
Observe the movement of fish everyday
Discourage entry of birds, goat, cow etc as they may carry germs (Since fish are susceptible to disease during winter months)

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	iyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अनुप - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION

Indian Council of Agricultural Research

Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Kiphire District

Bulletin No:100/2017

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 27th Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 22°C to 23°C and 11°C to 12°C, respectively. ▪ Relative humidity varied from 26% to 88%. ▪ Wind speed ranged from 3 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21°C - 22°C and the min temp 9°C to 10°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 21% to 75%. ▪ Wind speed may reach upto 2-4 kmph ▪ Wind direction will be southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties

Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
<i>Vegetables</i>			<i>Intercultural operations in vegetables are advised to remove the weeds.</i>

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अनु - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Kohima District

Bulletin No:100/2017

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 19 °C to 20°C and 11°C to 13°C, respectively.
- Relative humidity varied from 37% to 90%.
- Wind speed ranged from 3 to 4 kmph

- NDVI for Nagaland

Weather forecast valid upto 27th Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 19°C - 20°C and the **min temp** 6°C to 8°C
- **Sky is likely to be mainly clear** the coming week
- **Relative Humidity** is likely to range from 35% to 74%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be mostly easterly

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties
Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this</i>
---------------	--	-------------	---

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अनुप - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION

Indian Council of Agricultural Research

Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Longleng District

Bulletin No:100/2017

<i>Weather summary of the preceding week</i>		<i>Weather forecast valid upto 27th Dec'17</i>	
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 22°C to 23°C and 10°C to 13°C, respectively. ▪ Relative humidity varied from 60% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21°C - 22°C and the min temp 9°C to 10°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 21% to 75%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind direction will be mostly southeasterly 	
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 			
Field crops			
<p>Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties</p> <p>Mulching is essential to the rabi crops.</p>			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>
Horticultural crop			
<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
<i>Vegetables</i>			<i>Intercultural operations in vegetables are advised to remove the weeds.</i>

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अनुप - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Mokokchung District

Bulletin No:100/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

IA

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 27th Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 24^oC to 25^oC and 14^oC to 15^oC, respectively. ▪ Relative humidity varied from 42% to 90%. ▪ Wind speed ranged from 3 to 5 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21^oC - 22^oC and the min temp 9^oC to 10^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 21% to 75%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind direction will be southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties
Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove</i>
---------------	--	-------------	--

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा कृ अनुसंधान - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Mon District

Bulletin No:100/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Council of Agricultural Research)

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 27th Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 23°C to 25°C and 12°C to 14°C, respectively. ▪ Relative humidity varied from 36% to 85% ▪ Wind speed ranged from 5 to 6 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21°C - 22°C and the min temp 9°C to 10°C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 36% to 74%. ▪ Wind speed may reach upto 5- 6 kmph ▪ Wind direction will be southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties

Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove</i>
---------------	--	-------------	--

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा कृ अनुसंधान - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17

Peren District

Bulletin No:100/2017

<i>Weather summary of the preceding week</i>		<i>Weather forecast valid upto 27th Dec'17</i>	
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 22°C to 24°C and 12°C to 14°C, respectively. ▪ Relative humidity varied from 39% to 90%. ▪ Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21°C - 22°C and the min temp 9°C to 10°C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 37% to 74%. ▪ Wind speed may reach upto 3-5 kmph ▪ Wind direction will be southeasterly 	
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 			
<p>Field crops</p> <p>Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties</p> <p>Mulching is essential to the rabi crops.</p>			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>
<p>Horticultural crop</p>			
<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left</i>

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17

Phek District

Bulletin No:100/2017

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 22°C to 23°C and 11°C to 12°C, respectively.
- Relative humidity varied from 30% to 90%.
- Wind speed ranged from 1 to 2 kmph

- NDVI for Nagaland

Weather forecast valid upto 27th Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 21°C - 22°C and the **min temp** 9°C to 10°C
- **Sky is likely to be mainly clear** the coming week
- **Relative Humidity** is likely to range from 40% to 74%.
- **Wind speed** may reach upto 2-3 kmph
Wind direction will be mostly southeasterly

Field crops

Toriam and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties
Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum paddy	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
TRC/WRC paddy	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
Toria	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
Rabi maize	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
Linseed/len til	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy</i>
---------------	--	-------------	--

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	vyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Tuensang District

Bulletin No:100/2017

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 19^oC to 20^oC and 9^oC to 10^oC, respectively.
- Relative humidity varied from 30% to 95%.
- Wind speed ranged from 3 to 4 kmph

Weather forecast valid upto 27th Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 17^oC - 18^oC and the **min temp** 4^oC to 5^oC
- **Sky is likely to be mainly clear** the coming week
- **Relative Humidity** is likely to range from 22% to 70%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be mostly southeasterly

- NDVI for Nagaland

Field crops

Toriam and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties

Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
---------------	--	-------------	---

Sc

Sl.
o

1

2

3

4

5

6

7

8

ICAR RESEARCH COMPLEX FOR NEH REGION

Indian Council of Agricultural Research)



Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Wokha District

Bulletin No:100/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

Indian Council of Agricultural Research

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 27th Dec'17</i>
--	---

<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 22°C to 23°C and 12°C to 13°C, respectively. ▪ Relative humidity varied from 43% to 90%. ▪ Wind speed ranged from 4 to 5 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21°C - 22°C and the min temp 9°C to 10°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 24% to 75%. ▪ Wind speed may reach upto 2- 4 kmph ▪ Wind direction will be southeasterly
---	---

<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	
---	--

Field crops

Toriam and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties
Mulching is essential to the rabi crops.


Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

Citrus		Fruit flies	Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this
---------------	--	-------------	--

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





Integrated Agromet Advisory Service Bulletin from 23rd to 27th Dec'17 Zunheboto District

Bulletin No:100/2017

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 18°C to 19°C and 9°C to 10°C, respectively.
- Relative humidity varied from 37% to 85%.
- Wind speed ranged from 4 to 5 kmph

Weather forecast valid upto 27th Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 17°C - 18°C and the **min temp** 4°C to 6°C
- **Sky is likely to be mainly clear** the coming week
- **Relative Humidity** is likely to range from 22% to 70%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be mostly southeasterly

- NDVI for Nagaland

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties

Mulching is essential to the rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Harvest all mature panicle to reduce bird damage. Harvesting should be done on sunny day at least 15 cm above the ground for moisture conservation and Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles</i>
<i>Toria</i>	Vegetative stage	-	<i>Thinning and weeding is recommended in timely sown mustard crop Keep the field free from weeds and daily scouting of the field should be done for aphid</i>
<i>Rabi maize</i>	Vegetative stage		<i>Thinning must be done where more population was observed. Weeding and earthing up should be carried out</i>
<i>Linseed/lentil</i>	Sowing		<i>Continue sowing of lentil, linseed in rice field under zero tillage. Open a narrow furrow in between two rows of rice crop and sow the seeds in the furrow.</i>

Horticultural crop

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com

