



Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Dimapur District

Bulletin No:93/2017

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 29^oC to 30^oC and 15^oC to 17^oC, respectively. ▪ Relative humidity varied from 40% to 88%. ▪ Wind speed ranged from 3 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 25^oC - 26^oC and the min temp 14^oC to 15^oC ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 34% to 88%. ▪ Wind speed may reach upto 3- 4 kmph ▪ Wind direction will be mostly easterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toriam and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		<i>Provide mulching to conserve moisture</i>

<i>Horticultural crop</i>			
Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.			
<i>Citrus</i>		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
<i>Vegetables</i>	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>
<i>Cauliflower, cabbage,</i>			<i>Transplanting of cabbage and cauliflower should be done at a spacing – 30 x 60cm for small varieties. 45 x 60 cm for large varieties Keep the field free from weed. Hand hoeing should be done after transplanting</i>
<i>Litchi</i>			<i>Mulch the tree basin, protect young litchi plants from cold and suppress new flush by application of growth inhibitors</i>
<i>Ginger / turmeric</i>	Maturity stage		<i>Field must be inspected daily for disease appearance. While inspecting, the healthy plants/plots must be marked and kept for planting in the next season. Marking has to be done properly otherwise when plant matures and gets dried up, it will be very difficult to find out the marked plots.</i>
<i>Livestock</i>			
Vaccination of animals should be done against out-break of major diseases. Provide clean and cool drinking water to the animals			
<i>Poultry</i>	<ol style="list-style-type: none"> 1. Protect the birds from cold draft by covering the exposed windward areas using gunny bags and other locally available materials 2. Deworm the birds once every 2 months 		
<i>Piggery</i>	<ol style="list-style-type: none"> 1. Protect the animals from cold draft using gunny bags and other locally available materials 2. Provide protein rich ration and ad libitum clean drinking water 		
<i>Fisheries</i>			



If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha.
If water depth is less than 1m harvest 50% of the stocked fish

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	iyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17

Kiphire District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

A

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 23°C to 25°C and 11°C to 13°C, respectively. ▪ Relative humidity varied from 26% to 88%. ▪ Wind speed ranged from 3 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 23°C - 24°C and the min temp 13°C to 14°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 17% to 80%. ▪ Wind speed may reach upto 2 kmph ▪ Wind direction will be southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अतुप - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Kohima District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 21 °C to 22°C and 14°C to 15°C, respectively.
- Relative humidity varied from 37% to 95%.
- Wind speed ranged from 3 to 4 kmph

- NDVI for Nagaland

Weather forecast valid upto 3rd Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 20°C - 21°C and the **min temp** 11°C to 12°C
- **Sky is likely to be clear** the coming week
- **Relative Humidity** is likely to range from 31% to 84%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be mostly easterly

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots</i>
---------------	--	-------------	---

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अनुप - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

MA

Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Longleng District

Bulletin No:93/2017

<i>Weather summary of the preceding week</i>		<i>Weather forecast valid upto 3rd Dec'17</i>	
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 26°C to 27°C and 15°C to 17°C, respectively. ▪ Relative humidity varied from 60% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 22°C - 23°C and the min temp 11°C to 13°C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 34% to 78%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind direction will be southeasterly 	
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 			
Field crops			
<p>Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.</p> <p>Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.</p> <p>Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.</p>			
Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		<i>Provide mulching to conserve moisture</i>
Horticultural crop			
<p>Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.</p>			
Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17

Mokokchung District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

IA

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 24°C to 26°C and 14°C to 15°C, respectively. ▪ Relative humidity varied from 42% to 90%. ▪ Wind speed ranged from 3 to 5 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 25°C - 26°C and the min temp 14°C to 15°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 34% to 88%. ▪ Wind speed may reach upto 3-4 kmph ▪ Wind direction will be mostly easterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toriam and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
---------------	--	-------------	---

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा कृ अनुसंधान - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Mon District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 23°C to 254°C and 12°C to 14°C, respectively. ▪ Relative humidity varied from 36% to 95% ▪ Wind speed ranged from 5 to 6 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 25°C - 26°C and the min temp 14°C to 15°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 34% to 78%. ▪ Wind speed may reach upto 2- 4 kmph ▪ Wind direction will be easterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than</i>
---------------	--	-------------	--

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	iyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17

Peren District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

A

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 22°C to 24°C and 12°C to 14°C, respectively. ▪ Relative humidity varied from 39% to 95%. ▪ Wind speed ranged from 3 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 23°C - 24°C and the min temp 13°C to 14°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 17% to 80%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind direction will be south easterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left</i>
---------------	--	-------------	---

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा.कृ.अनुप. - पु.प.सं.
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Phek District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

Indian Council of Agricultural Research

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 22°C to 23°C and 11°C to 12°C, respectively. ▪ Relative humidity varied from 30% to 95%. ▪ Wind speed ranged from 1 to 2 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 23°C - 24°C and the min temp 12°C to 13°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 25% to 80%. ▪ Wind speed may reach upto 2-3kmph ▪ Wind direction will be mostly southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	vyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Tuensang District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

Sc

Sl.
o

1

2

3

4

5

6

7

8

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 19^oC to 20^oC and 9^oC to 10^oC, respectively. ▪ Relative humidity varied from 30% to 95%. ▪ Wind speed ranged from 3 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 18^oC - 20^oC and the min temp 9^oC to 10^oC ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 20% to 85%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind direction will be mostly southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.
Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the maggots may have already left the fruit to pupate</i>
Vegetables	Germination stage		<i>Thinning should be done and gap filling should be carried out</i>



Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Wokha District

Bulletin No:93/2017

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 3rd Dec'17</i>
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Maximum and minimum temperatures ranged 23°C to 24°C and 12°C to 13°C, respectively. ▪ Relative humidity varied from 43% to 92%. ▪ Wind speed ranged from 4 to 5 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 23°C - 24°C and the min temp 13°C to 15°C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 17% to 80%. ▪ Wind speed may reach upto 2- 4 kmph ▪ Wind direction will be mostly southeasterly
<ul style="list-style-type: none"> ▪ NDVI for Nagaland 	

Field crops

Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove and destroy any stung fruit, it is more effective to do this straight away than picking up rotten fruit from the ground as the</i>
---------------	--	-------------	---

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





Integrated Agromet Advisory Service Bulletin from 29th Nov to 3rd Dec'17 Zunheboto District

Bulletin No:93/2017

Weather summary of the preceding week

- No rain occurred the past week
- Maximum and minimum temperatures ranged 21°C to 22°C and 9°C to 11°C, respectively.
- Relative humidity varied from 37% to 95%.
- Wind speed ranged from 4 to 5 kmph

Weather forecast valid upto 3rd Dec'17

- Probability of no rain the coming week.
- **Max temp** is likely to be 19°C - 20°C and the **min temp** 9°C to 10°C
- **Sky is likely to be clear** the coming week
- **Relative Humidity** is likely to range from 29% to 84%.
- **Wind speed** may reach upto 4-5 kmph
- **Wind direction** will be mostly southeasterly

- NDVI for Nagaland

Field crops

Toriam and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.

Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles.

Mulching is essential to the rabi crops. Need based life saving irrigation is essential for rabi crops.

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum paddy</i>	Post Harvesting stage		<i>Follow sanitation during drying, milling and after milling to avoid contamination of grains and protect from insects, rodents and birds.</i>
<i>TRC/WRC paddy</i>	Dough development stage	Grasshopper	<i>It feeds irregularly on seedling and leaf blade and cuts the stem during the panicle stage. Bunds should be cleaned at regular interval.</i>
<i>Toria</i>	Sowing	-	<i>Quality seeds should be used. M-27, TS-36 and TS-38 are some of the varieties recommended for our region</i>
<i>Rabi maize</i>	Seedling stage		Provide mulching to conserve moisture

Horticultural crop

Use well drained soils, field sanitation should be maintained. Seed bed should be in good tilth, free from diseases and weeds.

Citrus		Fruit flies	<i>Once fruit is stung it is too late for anything but planning for a better result next year! You can check fruit for tiny dimples or weeping clear sap. Remove</i>
---------------	--	-------------	--

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com

