



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Dimapur	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 28^oC to 29^oC and 12^oC to 15^oC, respectively. ▪ Relative humidity varied from 36% to 80%. ▪ Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 27^oC - 28^oC and min temp 11^oC to 13^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 42% to 86%. ▪ Wind speed may reach upto 4-6 kmph ▪ Wind may blow from south easterly direction ▪ Weekly cumulative Rainfall: 0mm

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

- Intercultural operations in vegetables are advised to remove the weeds
- Marigold should be grown as border crop for chilli and tomato as a trap crop
- Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.
- Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done
<i>Toria/Mustard</i>	Vegetative stage	-	Daily monitoring should be done for pest and disease infestation
<i>Rabi maize</i>	Vegetative stage		Light hoeing should be done when necessary for better control of weeds
<i>Assam lemon</i>			Weeding should be done frequently. Daily monitoring should be done for aphid
Cabbage	Transplanted	Cabbage butterfly/caterpillar	In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation
Broccoli	Transplanted		In the present weather timely irrigation should be done
Radish, carrot	Transplanted		Timely earthing up should be done after weeding
Poultry	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide protein rich ration and ad libitum clean drinking water		
Fishery	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish		



“GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Kiphire	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26^oC to 27^oC and 9^oC to 10^oC, respectively. ▪ Relative humidity varied from 26% to 79%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 25^oC - 27^oC and min temp 8^oC to 10^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 24% to 84%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind may blow from easterly direction <p style="background-color: #ffffcc; margin-top: 5px;">Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Sky will be clear.

Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
Soybean	Post harvest stage		<i>It should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
Rabi maize	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Radish, turnip, Carrot	Transplanted		<i>In the present weather, maintain proper moisture condition in the soil. Never irrigate heavily as it will lead to excessive foliage, produce low quality roots and result in late maturity</i>
Pea, Rajmash	Vegetative stage		<i>Keeping weather in view, try to keep the plot weed free for proper growth and do timely earthing up</i>
Tomato	Transplanting stage	Leaf curl	<i>This disease is transmitted by white fly. Severe stunting of plants with downward rolling and crinkling of leaves, older leaves becomes brittle and thick. Try to apply carbofuran @ 1.5 kg /ha at the time of transplanting.</i>
Poultry	As the temperature is decreasing, the shed should be protected from cold wind by hanging curtain around the side of the wall or shed. If possible there should be separate space for different age group of birds. In the present weather, Sick bird should be immediately separated/ culled from healthy flock.		
Piggery	Observe their general stage of health, check for parasites and injuries, and watch out for any abnormal feeding habits		
Fishery	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Kohima	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 22⁰C to 23⁰C and 7⁰C to 9⁰C, respectively. ▪ Relative humidity varied from 42% to 97%. ▪ Wind speed ranged from 2 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 6⁰C to 7⁰C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 40% to 95%. ▪ Wind speed may reach upto 3-5 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0 mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
<i>Soybean</i>	Post Harvesting stage		<i>In the present weather, those that are harvested should be dried immediately</i>
<i>Toria/Mustard</i>	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Pea</i>	Germinating stage		<i>Keeping weather in view, mulching up should be carried out after earthing up</i>
<i>Tomato</i>	Transplanting stage		<i>Two rows of marigold for every 16 rows of tomato can be grown as a trap crop for pest management. Spray 5% neem seed kernel extract to kill early stages larvae.</i>
<i>Khasi mandarin</i>			<i>For good yield, maintain proper sanitation in the orchard</i>
<i>Winter vegetables</i>	Transplanting stage		<i>In the present weather transplanting of vegetables should be carried out.</i>
<i>Poultry</i>	Chickens and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks.		
<i>Piggery</i>	Provide high energy feeds like maize + wheat bran during cold days. Always disinfect the animal shed regularly.		
<i>Fishery</i>	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Longleng	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26⁰C to 27⁰C and 9⁰C to 11⁰C, respectively. ▪ Relative humidity varied from 43% to 80%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 25⁰C - 27⁰C and min temp 8⁰C to 10⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 39% to 82%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind may blow from south easterly direction

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		<i>Before storing of rice grains in storage, grains should be dried properly to bring down moisture level below 12%. Godowns should be thoroughly cleaned to destroy any insect-pests hiding in the creeks and crevices and dried before storing grains.</i>
Soybean	Post harvesting stage		<i>In the present weather, it should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
Rabi maize	Tasseling stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Khasi Mandarin	Fruiting to harvesting stage		<i>Mulching should be done with dry grasses near the tree base to conserve soil moisture</i>
Rajmash	Flowering stage		<i>In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Tomato	Transplanted		<i>In the present weather as transplanting is going on, provide irrigation after transplanting. Grow marigold, cowpea and beans as intercrop for better production.</i>
Poultry			<i>Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks</i>
Piggery			<i>As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days</i>
Fishery			<i>As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Mokokchung	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24⁰C to 26⁰C and 9⁰C to 10⁰C, respectively. ▪ Relative humidity varied from 41% to 86%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 26⁰C - 27⁰C and min temp 9⁰C to 10⁰C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 40% to 89%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		<i>Before storing of rice grains in storage, grains should be dried properly to bring down moisture level below 12%. Godowns should be thoroughly cleaned to destroy any insect-pests hiding in the creeks and crevices and dried before storing grains.</i>
Soybean	Post harvesting stage		<i>In the present weather, it should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
Rabi maize	Tasseling stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Khasi Mandarin	Fruiting to harvesting stage		<i>Mulching should be done with dry grasses near the tree base to conserve soil moisture</i>
Rajmash	Flowering stage		<i>In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Tomato	Transplanted		<i>In the present weather as transplanting is going on, provide irrigation after transplanting. Grow marigold, cowpea and beans as intercrop for better production.</i>
Poultry			<i>Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks</i>
Piggery			<i>As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days</i>
Fishery			<i>As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Mon	Bulletin No: 96/2018
Weather of preceding week	Weather forecast valid upto 05.12.2018	
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24^oC to 25^oC and 8^oC to 9^oC, respectively. ▪ Relative humidity varied from 45% to 92%. ▪ Wind speed ranged from 2 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 23^oC - 24^oC and min temp 6^oC to 7^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 45% to 92%. ▪ Wind speed may reach upto 2- 4 kmph ▪ Wind may blow from south easterly direction <p style="background-color: #ffffcc; margin-top: 5px;">Weekly cumulative Rainfall: 0mm</p>	

WEATHER FORECAST BASED AGROMET ADVISORIES
General Advisories: Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		<i>Before storing of rice grains in storage, grains should be dried properly to bring down moisture level below 12%. Godowns should be thoroughly cleaned to destroy any insect-pests hiding in the creeks and crevices and dried before storing grains.</i>
Soybean	Post harvest stage		<i>In the present weather proper drying should be done</i>
Potato	Sowing stage		<i>Sowing of potato should be done as soon as possible. Seed treatment with 2 gm Mancozeb and 1 gm Carbendazim mixed per litre of water per kilogram of seed for five minute should be done. Thereafter, seeds should be dried in shade before planting in the field</i>
Toria/Mustard	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
Pea	Sowing		<i>Keeping view in weather, sowing of pea should not be delayed, because late sowing will reduces seed yield and will exposes the crop to insect damage.</i>
Rabi maize	Vegetative stage		<i>Plants should be earthened up after every weeding.</i>
Citrus		Trunk borer	<i>For good yield , maintain proper sanitation in the orchard</i>
Radish	Vegetative stage		<i>In the present condition, earthing up should be done for proper development of roots. Radish has a tendency to bulge out of soil as it grows in size. Therefore, thorough covering by earthing up is recommended to produce quality roots</i>
Rajmash	Flowering stage		<i>In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any</i>
Poultry			<i>Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks Provide vitamin complex during sudden drop of temperature</i>
Piggery			<i>Get ready with provision of warmth for piglets during cold days in the upcoming winter with incandescent bulbs and wind breaks made from sack or plastic sheets</i>
Fishery			<i>If the water level is very low, partially harvest some of the stocked fishes</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Peren	Bulletin No: 96/2018
Weather of preceding week	Weather forecast valid upto 05.12.2018	
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24^oC to 25^oC and 8^oC to 9^oC, respectively. ▪ Relative humidity varied from 35% to 90%. ▪ Wind speed ranged from 3 to 4 kmph 	<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 24^oC - 24^oC and min temp 7^oC to 8^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 37% to 92%. ▪ Wind speed may reach upto 2- 4 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>	

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

- Intercultural operations in vegetables are advised to remove the weeds
- Marigold should be grown as border crop for chilli and tomato as a trap crop
- Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.
- Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done
<i>Toria/Mustard</i>	Vegetative stage	-	Daily monitoring should be done for pest and disease infestation
<i>Rabi maize</i>	Vegetative stage		Light hoeing should be done when necessary for better control of weeds
<i>Assam lemon</i>			Weeding should be done frequently. Daily monitoring should be done for aphid
Cabbage	Transplanted	Cabbage butterfly/caterpillar	In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation
Broccoli	Transplanted		In the present weather timely irrigation should be done
Radish, carrot	Transplanted		Timely earthing up should be done after weeding
Poultry	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide protein rich ration and ad libitum clean drinking water		
Fishery	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Phek	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 21⁰C to 22⁰C and 8⁰C to 9⁰C, respectively. ▪ Relative humidity varied from 32% to 92%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 5⁰C to 7⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 28% to 95%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from mostly south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
<i>Soybean</i>	Post Harvesting stage		<i>In the present weather, those that are harvested should be dried immediately</i>
<i>Toria/Mustard</i>	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Pea</i>	Germinating stage		<i>Keeping weather in view, mulching up should be carried out after earthing up</i>
<i>Tomato</i>	Transplanting stage		<i>Two rows of marigold for every 16 rows of tomato can be grown as a trap crop for pest management. Spray 5% neem seed kernel extract to kill early stages larvae.</i>
<i>Khasi mandarin</i>			<i>For good yield , maintain proper sanitation in the orchard</i>
<i>Winter vegetables</i>	Transplanting stage		<i>In the present weather transplanting of vegetables should be carried out.</i>
<i>Poultry</i>	Chickens and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks.		
<i>Piggery</i>	Provide high energy feeds like maize + wheat bran during cold days. Always disinfect the animal shed regularly.		
<i>Fishery</i>	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Tuensang	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 19⁰C to 20⁰C and 3⁰C to 4⁰C, respectively. ▪ Relative humidity varied from 18% to 84%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 18⁰C - 19⁰C and min temp 2⁰C to 3⁰C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 26% to 94%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Sky will be clear.

Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Ripening stage		<i>As the crop will be harvested by the coming week keep the threshing floor ready.</i>
Soybean	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
Rabi maize	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Radish, turnip, Carrot	Transplanted		<i>In the present weather, maintain proper moisture condition in the soil. Never irrigate heavily as it will lead to excessive foliage, produce low quality roots and result in late maturity</i>
Pea, Rajmash	Vegetative stage		<i>Keeping weather in view, try to keep the plot weed free for proper growth and do timely earthing up</i>
Tomato	Transplanting stage	Leaf curl	<i>This disease is transmitted by white fly. Severe stunting of plants with downward rolling and crinkling of leaves, older leaves becomes brittle and thick. Try to apply carbofuran @ 1.5 kg /ha at the time of transplanting.</i>
Poultry	As the temperature is decreasing, the shed should be protected from cold wind by hanging curtain around the side of the wall or shed. If possible there should be separate space for different age group of birds. In the present weather, Sick bird should be immediately separated/ culled from healthy flock.		
Piggery	Observe their general stage of health, check for parasites and injuries, and watch out for any abnormal feeding habits		
Fishery	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Wokha	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24^oC to 25^oC and 8^oC to 10^oC respectively. ▪ Relative humidity varied from 50% to 89%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 23^oC - 24^oC and min temp 7^oC to 8^oC ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 46% to 95%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		<i>Before storing of rice grains in storage, grains should be dried properly to bring down moisture level below 12%. Godowns should be thoroughly cleaned to destroy any insect-pests hiding in the creeks and crevices and dried before storing grains.</i>
Soybean	Post harvesting stage		<i>In the present weather, it should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
Rabi maize	Tasseling stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Khasi Mandarin	Fruiting to harvesting stage		<i>Mulching should be done with dry grasses near the tree base to conserve soil moisture</i>
Rajmash	Flowering stage		<i>In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Tomato	Transplanted		<i>In the present weather as transplanting is going on, provide irrigation after transplanting. Grow marigold, cowpea and beans as intercrop for better production.</i>
Poultry			<i>Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks</i>
Piggery			<i>As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days</i>
Fishery			<i>As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface</i>





GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 30.11.18	District: Zunheboto	Bulletin No: 96/2018
Weather of preceding week		Weather forecast valid upto 05.12.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24^oC to 26^oC and 8^oC to 10^oC, respectively. ▪ Relative humidity varied from 31% to 90%. ▪ Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 23^oC - 24^oC and min temp 6^oC to 8^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 34% to 95%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Intercultural operations in vegetables are advised to remove the weeds

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Storage		<i>Before storing of rice grains in storage, grains should be dried properly to bring down moisture level below 12%. Godowns should be thoroughly cleaned to destroy any insect-pests hiding in the creeks and crevices and dried before storing grains.</i>
Soybean	Post harvesting stage		<i>In the present weather, it should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	Aphids	<i>In the present weather proper monitoring should be done for aphids</i>
Rabi maize	Tasseling stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Khasi Mandarin	Fruiting to harvesting stage		<i>Mulching should be done with dry grasses near the tree base to conserve soil moisture</i>
Rajmash	Flowering stage		<i>In the present weather condition, proper mulching should be done. Collect and destroy disease infected and insect infested plant parts if there is any</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Tomato	Transplanted		<i>In the present weather as transplanting is going on, provide irrigation after transplanting. Grow marigold, cowpea and beans as intercrop for better production.</i>
Poultry			<i>Chicks and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks</i>
Piggery			<i>As the days are getting cooler, provide high energy feeds like maize + wheat bran during cold days</i>
Fishery			<i>As far as possible do not disturb the pond. Prune the tree branches along the pond embankment and allow enough sunlight to fall on the pond water surface</i>

