



Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Dimapur District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 30^oC to 33^oC and 21^oC to 24^oC, respectively.
- Relative humidity varied from 35% to 90%.
- Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 20th June 2018


- Probability of moderate rain the coming week.
- **Max temp** is likely to be 30^oC - 31^oC and the **min temp** 24^oC to 25^oC
- **Sky is likely to be partly cloudy** the coming week
- **Relative Humidity** is likely to range from 47% to 90%.
- **Wind speed** may reach upto 2- 3 kmph
- **Wind direction** will be mostly southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum Paddy</i>	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
<i>Lowland paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Green gram</i>	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>

Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>
Horticultural crop			
<ul style="list-style-type: none"> For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations <i>Trichoderma harzianum</i> and <i>T. viridae</i> can be used for seed and soil treatment as well as foliar application against the fungal pathogens, Crop rotation should be followed for starving the pathogens and declining their population, One percent Bordeaux mixture can be applied as root dip, soil drench and foliar spray, bordeaux paint applied on the trunk and pruned wounds of citrus prevent pathogens attack Banana crop is infested by a number of pest and disease. Disease can be controlled with frequent applications of fungicides, certifying seed, and cultural practices, such as practise of good hygiene, avoid large areas of any one variety of crop, especially in high disease areas, regular monitor of crops for diseases and resistance, and adequate spacing of plants and efficient drainage within plantation 			
Khasi Mandarin	New flush		<i>Proper monitoring should be done for trunk borers. Collect and destroy them wherever feasible and inject petrol or nuvan into the holes and plug the holes with mud.</i>
Cucurbits	Vegetative to flowering stage	Red pumpkin beetle	<i>Monitor the plants against red pumpkin beetle in cucurbits especially bottle gourd, pumpkin, cucumber, water melon etc. Adults are small, elongated yellow and defoliate the leaves immediately after germination. Larvae feed on roots and plant parts. Mechanically collect and destroy the pest if incidence is low.</i>
Okra	Fruiting stage	Fruit and shoot borer	<i>Collect and destroy all the infested fruits. Timely earthing up should be done</i>
Brinjal	Ratoon crop	Shoot and fruit borer	<i>Incidence of Shoot and fruit borer usually occurs during humid conditions after the rainfall. Small brown caterpillars bore into the top tender shoots and tunnel downwards the main axis which wither, droop down and growing points are killed and later on they bore into the fruits and feed within. Affected fruits become unfit for consumption. Infested fruits and shoots should be removed regularly and buried deep in the soil.</i>
Ginger / turmeric	Vegetative stage	Rhizome rot	<i>The plot should be well drained free from water logging and while planting healthy rhizomes should be used for cultivation so as to prevent rhizome rot. Maintain proper drainage in the field of ginger to prevent diseases. The field must be inspected regularly for disease appearance and more often when it rains.</i>
Livestock			
<ul style="list-style-type: none"> Water stagnation should be avoided in livestock shed to avoid mosquito breeding 			
Poultry	<ul style="list-style-type: none"> Shades from tall trees and plantation around the sheds can reduce the radiant heat. The plantation of trees should be such that trees will be leafy during summer and bald during winter Thatching of roof with paddy straw or sugar cane leaves will reduce temperature inside the shed For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to 		

		<p>avoid dehydration after transportation</p> <ul style="list-style-type: none"> • Shifting, transportation, de-beaking and vaccination should be done during night or cool hours of the day
	Piggery	<ul style="list-style-type: none"> • Identify and isolate the infected and in contact animals • Dispose the dead animals either by burning or deep burial • Regular disinfection of shed and its premises with 1-2 % phenyl • Regular de-worming to control internal parasites • Keep the animal house clean and dry
	Fisheries	
		<ul style="list-style-type: none"> • By this month, renovation of old pond and excavation of new pond should be completed. • Proper inspection of work should be done for new pond • Adopt culture practice as per the technical advice <p>If the management measures are followed accordingly, a good harvest can be expected. It is advisable to consult a fisheries expert prior to taking any management measures</p>

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा कृ अनुप - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Kiphire District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 27°C to 29°C and 19°C to 21°C, respectively.
- Relative humidity varied from 53% to 93%.
- Wind speed ranged from 3 to 4 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 25°C - 26°C and the **min temp** 19°C to 20°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 47% to 95%.
- **Wind speed** may reach upto 1-2 kmph
- **Wind direction** will be mostly southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
TRC/WRC paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations *Trichoderma harzianum* and *T. viridae* can be used for seed and soil treatment as well as foliar application against the fungal

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Kohima District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 24 °C to 26°C and 17°C to 19°C, respectively.
- Relative humidity varied from 60% to 95%.
- Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 22°C - 24°C and the **min temp** 16°C to 17°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 63% to 95%.
- **Wind speed** may reach upto 2-3 kmph
- **Wind direction** will be mostly southerly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
TRC/WRC paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा क अनुप - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Longleng District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 28°C to 30°C and 20°C to 21°C, respectively.
- Relative humidity varied from 54% to 94%.
- Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 25°C - 26°C and the **min temp** 19°C to 20°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 55% to 95%.
- **Wind speed** may reach upto 2- 3kmph
Wind direction will be mostly southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
TRC/WRC paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management, proper drainage and raised beds are a must for management of nursery diseases. formulations

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18

Mokokchung District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 27°C to 30°C and 19°C to 20°C, respectively.
- Relative humidity varied from 29% to 93%.
- Wind speed ranged from 3 to 5 kmph

Weather forecast valid upto 20th June 2018

- Probability of heavy rain the coming week.
- **Max temp** is likely to be 25°C - 27°C and the **min temp** 19°C to 20°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 64% to 95%.
- **Wind speed** may reach upto 2- 4 kmph
- **Wind direction** will be southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
TRC/WRC paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations *Trichoderma harzianum* and *T. viridae* can be used for seed and soil treatment as well as foliar application against the fungal pathogens. Crop rotation should be followed for starving the pathogens and declining their population. One percent

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



भा कृ अनुसंधान - पु प सं
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION

(Indian Council of Agricultural Research)

Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Mon District

Bulletin No:48/2018

<i>Weather summary of the preceding week</i>	<i>Weather forecast valid upto 20th June 2018</i>
<ul style="list-style-type: none"> ▪ Moderate rain occurred the past week ▪ Maximum and minimum temperatures ranged 25°C to 28°C and 19°C to 20°C, respectively. ▪ Relative humidity varied from 65% to 90% ▪ Wind speed ranged from 2 to 3 kmph 	<ul style="list-style-type: none"> ▪ Probability of moderate rain the coming week. ▪ Max temp is likely to be 24°C - 25°C and the min temp 18°C to 19°C ▪ Sky is likely to be cloudy the coming week ▪ Relative Humidity is likely to range from 60% to 93%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind direction will be mostly southeasterly
<ul style="list-style-type: none"> • Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable. • Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well. • Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer. 	

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum Paddy</i>	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Green gram</i>	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
<i>Maize</i>	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations *Trichoderma harzianum* and *T. viridae* can be used for seed and soil treatment as well as foliar application against the fungal pathogens, Crop rotation should be followed for starving the pathogens and declining their population, One percent Bordeaux mixture can be applied as root dip, soil drench and foliar spray, bordeaux paint applied on the trunk and pruned wounds of citrus prevent pathogens attack
- Banana crop is infested by a number of pest and disease. Disease can be controlled with frequent applications of fungicides, certifying seed, and cultural practices, such as practise of good hygiene, avoid large areas of any one variety of crop, especially in high disease areas, regular monitor of crops for diseases and resistance, and adequate spacing of plants and efficient drainage within plantation

<i>Khasi Mandarin</i>	New flush		<i>Proper monitoring should be done for trunk borers. Collect and destroy them wherever feasible and inject petrol or nuvan into the holes and plug the holes with mud.</i>
<i>Cucurbits</i>	Vegetative to flowering stage	Red pumpkin beetle	<i>Monitor the plants against red pumpkin beetle in cucurbits especially bottle gourd, pumpkin, cucumber, water melon etc. Adults are small, elongated yellow and defoliate the leaves immediately after germination. Larvae feed on roots and plant parts. Mechanically collect and destroy the pest if incidence is low.</i>
<i>Okra</i>	Fruiting stage	Fruit and shoot borer	<i>Collect and destroy all the infested fruits. Timely earthing up should be done</i>
<i>Brinjal</i>	Ratoon crop	Shoot and fruit borer	<i>Incidence of Shoot and fruit borer usually occurs during humid conditions after the rainfall. Small brown caterpillars bore into the top tender shoots and tunnel downwards the main axis which wither, droop down and growing points are killed and later on they bore into the fruits and feed within. Affected fruits become unfit for consumption. Infested fruits and shoots should be removed regularly and buried deep in the soil.</i>
<i>Ginger / turmeric</i>	Vegetative stage	Rhizome rot	<i>The plot should be well drained free from water logging and while planting healthy rhizomes should be used for cultivation so as to prevent rhizome rot. Maintain proper drainage in the field of ginger to prevent diseases. The field must be inspected regularly for disease appearance and more often when it rains.</i>

Livestock

- Water stagnation should be avoided in livestock shed to avoid mosquito breeding

<i>Poultry</i>	<ul style="list-style-type: none"> • Shades from tall trees and plantation around the sheds can reduce the radiant heat. The plantation of trees should be such that trees will be leafy during summer and bald during winter • Thatching of roof with paddy straw or sugar cane leaves will reduce temperature inside the shed • For day old chicks provide cool water and electrolytes on their arrival to farm before offering feed to avoid dehydration after transportation • Shifting, transportation, de-beaking and vaccination should be done during night or cool hours of the day
<i>Piggery</i>	<ul style="list-style-type: none"> • Identify and isolate the infected and in contact animals • Dispose the dead animals either by burning or deep burial • Regular disinfection of shed and its premises with 1-2 % phenyl • Regular de-worming to control internal parasites • Keep the animal house clean and dry

Fisheries



भा कृ अनुसंधान - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Peren District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 26°C to 27°C and 19°C to 20°C, respectively.
- Relative humidity varied from 60% to 94%.
- Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 23°C - 25°C and the **min temp** 18°C to 19°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 73% to 96%.
- **Wind speed** may reach upto 2-3 kmph
Wind direction will be mostly southerly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
Lowland paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Phek District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 25°C to 27°C and 18°C to 19°C, respectively.
- Relative humidity varied from 28% to 92%.
- Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 24°C - 26°C and the **min temp** 17°C to 18°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 59% to 95%.
- **Wind speed** may reach upto 2-3 kmph
- **Wind direction** will be mostly southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
TRC/WRC paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	vyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Tuensang District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 25^oC to 28^oC and 14^oC to 16^oC, respectively.
- Relative humidity varied from 65% to 95%.
- Wind speed ranged from 2 to 3 kmph

Weather forecast valid upto 20th June 2018

- Probability of heavy rain the coming week.
- **Max temp** is likely to be 21^oC - 25^oC and the **min temp** 13^oC to 15^oC
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 56% to 95%.
- **Wind speed** may reach upto 2-3 kmph
- **Wind direction** will be mostly southerly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum Paddy</i>	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Green gram</i>	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
<i>Maize</i>	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations *Trichoderma harzianum* and *T. viridae* can be used for seed and soil treatment as well as foliar application against the fungal pathogens, Crop rotation should be followed for starving the pathogens and declining their population, One percent Bordeaux mixture can be applied as root dip, soil drench and foliar spray, bordeaux paint applied on the trunk

Sc

Sl.
o

1

2

3

4

5

6

7

8

ICAR RESEARCH COMPLEX FOR NEH REGION

Horticultural Research)



Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18

Wokha District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 26^oC to 29^oC and 19^oC to 20^oC, respectively.
- Relative humidity varied from 60% to 91%.
- Wind speed ranged from 3 to 4 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 22^oC - 24^oC and the **min temp** 18^oC to 19^oC
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 82% to 92%.
- **Wind speed** may reach upto 2- 3 kmph
- **Wind direction** will be southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.


Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum Paddy</i>	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
<i>TRC/WRC paddy</i>	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
<i>Green gram</i>	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
<i>Maize</i>	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





भा क अनुप - पु प सं
ICAR - RCNEH

Integrated Agromet Advisory Service Bulletin from 16th to 20th June '18 Zunheboto District

Bulletin No:48/2018

Weather summary of the preceding week

- Moderate rain occurred the past week
- Maximum and minimum temperatures ranged 25°C to 27°C and 18°C to 19°C, respectively.
- Relative humidity varied from 75% to 94%.
- Wind speed ranged from 2 to 4 kmph

Weather forecast valid upto 20th June 2018

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 23°C - 25°C and the **min temp** 18°C to 19°C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 69% to 95%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be southeasterly

- Locally available nutrient resources should always be preferred along with the external nutrient inputs to make the farming system more sustainable.
- Neem products should be incorporated as they are non-toxic and are a safe alternative to commercially manufactured chemical fertilisers. Spraying should be undertaken in the morning or late in the afternoon. Insects lay eggs on the underside of the leaves. Hence it is important to spray on the underside of the leaves as well.
- Low cost rain water harvesting structure (Jalkund) should be incorporated as it provides irrigation in lean season and provides healthy food. It also increases the production and productivity of high value vegetables crops and generates additional net monetary income of the farmer.

Field crops

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
Jhum Paddy	Tillering stage	-	<i>High humidity favour the disease incidence. It is a severe fungal disease. Typical spindle shaped leaf lesions, wide at centre and pointed toward either end are seen. In severe cases they coalesce and the leaves die. Neck and panicle infection causes chaffy and shriveled grains. Maintain rice fields free from weeds which give shelter to many pathogens</i>
TRC/WRC paddy	Nursery		<i>Prepare the nursery by plowing at least twice and harrow at least once. Level the soil surface and put in drainage lines across the field. Broadcast seed in the nursery evenly, over the water covered soil surface Seed treatment should be done</i>
Green gram	Podding stage	-	<i>Clean cultivation should be maintained. Proper earthing up should be done</i>
Maize	Silking stage		<i>Timely weeding and earthing up should be done so as to prevent the plants from lodging</i>

Horticultural crop

- For disease management proper drainage and raised beds are a must for management of nursery diseases, formulations

Scientific Expert committee

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com

