



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18

## Dimapur District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>0</sup>C to 34<sup>0</sup>C and 17<sup>0</sup>C to 20<sup>0</sup>C, respectively.
- Relative humidity varied from 42% to 96%.
- Wind speed ranged from 1 to 2 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 30<sup>0</sup>C - 31<sup>0</sup>C and the **min temp** 19<sup>0</sup>C to 20<sup>0</sup>C
- **Sky is likely to be partly cloudy** the coming week
- **Relative Humidity** is likely to range from 26% to 87%.
- **Wind speed** may reach upto 2- 4 kmph
- **Wind direction** will be mostly easterly

- Timely weeding and earthing up should be carried out


### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	Clean cultivation should be maintained
<i>Maize</i>	vegetative stage		Timely weeding and earthing up should be done for the plant to grow well
<i>Water harvesting</i>			Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
<i>Cucurbits</i>	Vegetative stage		<i>Provide Proper staking</i>
<i>Okra</i>	Flowering stage	-	<i>Timely earthing up should be done</i>

	<b>Brinjal</b>	Field preparation	<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	- <i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
<b>Livestock</b>			
<ul style="list-style-type: none"> <li>• Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>			
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>• Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>• In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>• Birds must be vaccinated against infectious diseases</li> <li>• Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>	
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>• Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>• Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>	
<b>Fisheries</b>			
 <p>भारतीय ICAR</p>		<ul style="list-style-type: none"> <li>• Lime pond as per recommended dose @10kg/bigha</li> <li>• Stock fish fingerling @8000/ha</li> <li>• Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>• If possible, stock bigger size fish seed to ensure faster growth</li> </ul>	

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18

## Kiphire District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>o</sup>C to 28<sup>o</sup>C and 15<sup>o</sup>C to 16<sup>o</sup>C, respectively.
- Relative humidity varied from 23% to 85%.
- Wind speed ranged from 3 to 4 kmph


### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 25<sup>o</sup>C - 26<sup>o</sup>C and the **min temp** 15<sup>o</sup>C to 16<sup>o</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 29% to 90%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be mostly easterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too

### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	<i>Clean cultivation should be maintained</i>
<i>Maize</i>	vegetative stage		<i>Timely weeding and earthing up should be done for the plant to grow well</i>
<i>Water harvesting</i>			<i>Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price</i>
<b><i>Horticultural crop</i></b>			
<ul style="list-style-type: none"> <li>• Standing rain water in the field may be drained off immediately after rain</li> <li>• Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping</li> <li>• Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting</li> <li>• Transplanting of King chili should be done</li> </ul>			
<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
<i>Cucurbits</i>	Vegetative stage		<i>Provide Proper staking</i>
<i>Okra</i>	Flowering stage	-	<i>Timely earthing up should be done</i>

<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing <i>Pseudomonas aeruginosa</i>, <i>Trichoderma harzianum</i> and <i>Bacillus brevis</i> for 1 hr.</i>
<b>Livestock</b>			
<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>			
<b>Poultry</b>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
<b>Piggery</b>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b>Fisheries</b>			
	<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @ 10kg/bigha</li> <li>Stock fish fingerling @ 8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May'18

## Kohima District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 23<sup>o</sup>C to 26<sup>o</sup>C and 14<sup>o</sup>C to 15<sup>o</sup>C, respectively.
- Relative humidity varied from 34% to 90%.
- Wind speed ranged from 3 to 4 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of moderate rain the coming week.
- **Max temp** is likely to be 22<sup>o</sup>C - 23<sup>o</sup>C and the **min temp** 13<sup>o</sup>C to 14<sup>o</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 47% to 92%.
- **Wind speed** may reach upto 2-3 kmph
- **Wind direction** will be mostly easterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too


### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	<i>Clean cultivation should be maintained</i>
<i>Maize</i>	vegetative stage		<i>Timely weeding and earthing up should be done for the plant to grow well</i>
<i>Water harvesting</i>			<i>Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price</i>

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
-----------------------	--	--	--

	<b>Cucurbits</b>	Vegetative stage		<i>Provide Proper staking</i>
	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
	<b>Khasi Mandarin</b>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
<b>Livestock</b>				
<ul style="list-style-type: none"> <li>• Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>				
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>• Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>• In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>• Birds must be vaccinated against infectious diseases</li> <li>• Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>• Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>• Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b>Fisheries</b>				
		<ul style="list-style-type: none"> <li>• Lime pond as per recommended dose @10kg/bigha</li> <li>• Stock fish fingerling @8000/ha</li> <li>• Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>• If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		
 <p>भारत सरकार ICAR</p>				



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com




भा क अनुष - पु प सं  
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION  
(Indian Council of Agricultural Research)  
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

## Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18 Longleng District

**Bulletin No:35/2018**

<i>Weather summary of the preceding week</i>			<i>Weather forecast valid upto 6<sup>th</sup> May 2018</i>
<ul style="list-style-type: none"> <li>Light rain occurred the past week</li> <li>Maximum and minimum temperatures ranged 24<sup>0</sup>C to 27<sup>0</sup>C and 17<sup>0</sup>C to 18<sup>0</sup>C, respectively.</li> <li>Relative humidity varied from 23% to 85%.</li> <li>Wind speed ranged from 3 to 4 kmph</li> </ul>			<ul style="list-style-type: none"> <li>Probability of light rain the coming week.</li> <li><b>Max temp</b> is likely to be 25<sup>0</sup>C - 26<sup>0</sup>C and the <b>min temp</b> 15<sup>0</sup>C to 16<sup>0</sup>C</li> <li><b>Sky is likely to be mainly cloudy</b> the coming week</li> <li><b>Relative Humidity</b> is likely to range from 31% to 89%.</li> <li><b>Wind speed</b> may reach upto 2- 3 kmph</li> <li><b>Wind direction</b> will be mostly southeasterly</li> </ul>
<ul style="list-style-type: none"> <li>Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too</li> </ul>			
<i>Field crops</i>			
<b>Main Crops</b>	<b>Stage</b>	<b>Pest/ Diseases</b>	<b>Agro-meteorological Advisories</b>
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>Place sticky traps in the drying and storage areas for rats.</li> <li>Storage rooms should be physically rodent and bird proof, if possible.</li> <li>Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	Clean cultivation should be maintained
<i>Maize</i>	vegetative stage		Timely weeding and earthing up should be done for the plant to grow well
<i>Water harvesting</i>			Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price
<i>Horticultural crop</i>			
<ul style="list-style-type: none"> <li>Standing rain water in the field may be drained off immediately after rain</li> <li>Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping</li> <li>Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting</li> <li>Transplanting of King chili should be done</li> </ul>			
<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer
<i>Cucurbits</i>	Vegetative stage		Provide Proper staking

	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
<b>Livestock</b>				
<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>				
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b>Fisheries</b>				
		<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @ 10kg/bigha</li> <li>Stock fish fingerling @ 8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18

## Mokokchung District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 28<sup>o</sup>C to 30<sup>o</sup>C and 14<sup>o</sup>C to 15<sup>o</sup>C, respectively.
- Relative humidity varied from 29% to 80%.
- Wind speed ranged from 3 to 5 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 25<sup>o</sup>C - 26<sup>o</sup>C and the **min temp** 15<sup>o</sup>C to 16<sup>o</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 45% to 90%.
- **Wind speed** may reach upto 3 - 4 kmph
- **Wind direction** will be southeasterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too

### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	<i>Clean cultivation should be maintained</i>
<i>Maize</i>	vegetative stage		<i>Timely weeding and earthing up should be done for the plant to grow well</i>
<i>Water harvesting</i>			<i>Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price</i>

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
<i>Cucurbits</i>	Vegetative stage		<i>Provide Proper staking</i>

<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing <i>Pseudomonas aeruginosa</i>, <i>Trichoderma harzianum</i> and <i>Bacillus brevis</i> for 1 hr.</i>
<b><i>Livestock</i></b>			
<ul style="list-style-type: none"> <li>• Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>			
<b>Poultry</b>	<ul style="list-style-type: none"> <li>• Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>• In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>• Birds must be vaccinated against infectious diseases</li> <li>• Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
<b>Piggery</b>	<ul style="list-style-type: none"> <li>• Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>• Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b><i>Fisheries</i></b>			
	<ul style="list-style-type: none"> <li>• Lime pond as per recommended dose @10kg/bigha</li> <li>• Stock fish fingerling @8000/ha</li> <li>• Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>• If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18

## Mon District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>o</sup>C to 28<sup>o</sup>C and 14<sup>o</sup>C to 16<sup>o</sup>C, respectively.
- Relative humidity varied from 33% to 87%
- Wind speed ranged from 3 to 5 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 24<sup>o</sup>C - 25<sup>o</sup>C and the **min temp** 14<sup>o</sup>C to 15<sup>o</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 39% to 90%.
- **Wind speed** may reach upto 2- 4 kmph
- **Wind direction** will be mostly southeasterly

Timely weeding and earthing up should be carried out

### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Jhum Paddy</i>			<ul style="list-style-type: none"> <li>• Maize and pulses should be grown as strip crop in hill slopes</li> <li>• Use certified seeds and if there is a history of occurrence of diseases in that particular field then use a resistant variety.</li> </ul>
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	Clean cultivation should be maintained
<i>Maize</i>	vegetative stage		Timely weeding and earthing up should be done for the plant to grow well
<i>Water harvesting</i>			Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer
-----------------------	--	--	---



	<b>Cucurbits</b>	Vegetative stage		<i>Provide Proper staking</i>
	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
<b>Livestock</b>				
<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>				
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b>Fisheries</b>				
		<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @10kg/bigha</li> <li>Stock fish fingerling @8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18

## Peren District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>o</sup>C to 29<sup>o</sup>C and 14<sup>o</sup>C to 15<sup>o</sup>C, respectively.
- Relative humidity varied from 30% to 87%.
- Wind speed ranged from 3 to 4 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 24<sup>o</sup>C - 25<sup>o</sup>C and the **min temp** 14<sup>o</sup>C to 15<sup>o</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 46% to 95%.
- **Wind speed** may reach upto 2-3 kmph
- **Wind direction** will be mostly southeasterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too

### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	<i>Clean cultivation should be maintained</i>
<i>Maize</i>	vegetative stage		<i>Timely weeding and earthing up should be done for the plant to grow well</i>
<i>Water harvesting</i>			<i>Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price</i>

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
-----------------------	--	----------------------------------	--

			borer	
	<b>Cucurbits</b>	Vegetative stage		<i>Provide Proper staking</i>
	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
	<b>Livestock</b>			
	<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>			
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
	<b>Fisheries</b>			
		<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @ 10kg/bigha</li> <li>Stock fish fingerling @ 8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18 Phek District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 23<sup>0</sup>C to 24<sup>0</sup>C and 14<sup>0</sup>C to 15<sup>0</sup>C, respectively.
- Relative humidity varied from 28% to 85%.
- Wind speed ranged from 3 to 4 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 23<sup>0</sup>C - 24<sup>0</sup>C and the **min temp** 14<sup>0</sup>C to 15<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 35% to 90%.
- **Wind speed** may reach upto 2-3 kmph  
**Wind direction** will be southeasterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too


### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	Clean cultivation should be maintained
<i>Maize</i>	vegetative stage		Timely weeding and earthing up should be done for the plant to grow well
<i>Water harvesting</i>			Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk	Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer
-----------------------	--	----------------------------------	---

			borer	
	<i>Cucurbits</i>	Vegetative stage		<i>Provide Proper staking</i>
	<i>Okra</i>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<i>Brinjal</i>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<i>Ginger / turmeric</i>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
<b><i>Livestock</i></b>				
<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>				
	<i>Poultry</i>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<i>Piggery</i>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b><i>Fisheries</i></b>				
		<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @10kg/bigha</li> <li>Stock fish fingerling @8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		

<b>Sl.no</b>	<b>Name</b>	<b>Designation</b>	<b>Department</b>	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18 Tuensang District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 25<sup>0</sup>C to 27<sup>0</sup>C and 13<sup>0</sup>C to 14<sup>0</sup>C, respectively.
- Relative humidity varied from 29% to 80%.
- Wind speed ranged from 3 to 4 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*


- Probability of light rain the coming week.
- **Max temp** is likely to be 22<sup>0</sup>C - 23<sup>0</sup>C and the **min temp** 12<sup>0</sup>C to 13<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 36% to 95%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be southeasterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too

### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	<i>Clean cultivation should be maintained</i>
<i>Maize</i>	vegetative stage		<i>Timely weeding and earthing up should be done for the plant to grow well</i>
<i>Water harvesting</i>			<i>Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price</i>
<b><i>Horticultural crop</i></b>			
<ul style="list-style-type: none"> <li>• Standing rain water in the field may be drained off immediately after rain</li> <li>• Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping</li> <li>• Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting</li> <li>• Transplanting of King chili should be done</li> </ul>			
<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>

ICAR RESEARCH COMPLEX FOR NEH REGION  
 (Indian Council of Agricultural Research)  
 NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

	<b>Cucurbits</b>	Vegetative stage		<i>Provide Proper staking</i>
	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
	<b>Livestock</b>			
<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>				
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b>Fisheries</b>				
		<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @10kg/bigha</li> <li>Stock fish fingerling @8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		

<b>Sl.no</b>	<b>Name</b>	<b>Designation</b>	<b>Department</b>	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com



# Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18

## Wokha District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 26<sup>o</sup>C to 29<sup>o</sup>C and 14<sup>o</sup>C to 15<sup>o</sup>C, respectively.
- Relative humidity varied from 32% to 85%.
- Wind speed ranged from 4 to 5 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 24<sup>o</sup>C - 25<sup>o</sup>C and the **min temp** 14<sup>o</sup>C to 15<sup>o</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 52% to 91%.
- **Wind speed** may reach upto 3- 4 kmph
- **Wind direction** will be southeasterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too

### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	Clean cultivation should be maintained
<i>Maize</i>	vegetative stage		Timely weeding and earthing up should be done for the plant to grow well
<i>Water harvesting</i>			Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk borer	<i>Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer</i>
-----------------------	--	--	--

	<b>Cucurbits</b>	Vegetative stage		<i>Provide Proper staking</i>
	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
	<b>Livestock</b>			
	<ul style="list-style-type: none"> <li>• Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>			
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>• Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>• In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>• Birds must be vaccinated against infectious diseases</li> <li>• Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>• Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>• Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
	<b>Fisheries</b>			
		<ul style="list-style-type: none"> <li>• Lime pond as per recommended dose @ 10kg/bigha</li> <li>• Stock fish fingerling @ 8000/ha</li> <li>• Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>• If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		



## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com





--	--





भा क अनुप - पु प सं  
ICAR - RCNEH

ICAR RESEARCH COMPLEX FOR NEH REGION  
(Indian Council of Agricultural Research)  
NAGALAND CENTER, JHARNAPANI, MEDZIPHEMA

## Integrated Agromet Advisory Service Bulletin from 2<sup>nd</sup> to 6<sup>th</sup> May '18 Zunheboto District

**Bulletin No:35/2018**

### *Weather summary of the preceding week*

- Light rain occurred the past week
- Maximum and minimum temperatures ranged 25<sup>0</sup>C to 29<sup>0</sup>C and 14<sup>0</sup>C to 15<sup>0</sup>C, respectively.
- Relative humidity varied from 31% to 84%.
- Wind speed ranged from 4 to 5 kmph

### *Weather forecast valid upto 6<sup>th</sup> May 2018*

- Probability of light rain the coming week.
- **Max temp** is likely to be 24<sup>0</sup>C - 25<sup>0</sup>C and the **min temp** 14<sup>0</sup>C to 15<sup>0</sup>C
- **Sky is likely to be mainly cloudy** the coming week
- **Relative Humidity** is likely to range from 43% to 90%.
- **Wind speed** may reach upto 3-4 kmph
- **Wind direction** will be southeasterly

- Timely weeding and earthing up should be carried out as there is probability of light rain the coming week too


### *Field crops*

Main Crops	Stage	Pest/ Diseases	Agro-meteorological Advisories
<i>Paddy</i>	Storage		<ul style="list-style-type: none"> <li>• Place sticky traps in the drying and storage areas for rats.</li> <li>• Storage rooms should be physically rodent and bird proof, if possible.</li> <li>• Inspect the stored seeds once a week for signs of insect infestation.</li> </ul>
<i>Green gram</i>	Vegetative stage	-	Clean cultivation should be maintained
<i>Maize</i>	vegetative stage		Timely weeding and earthing up should be done for the plant to grow well
<i>Water harvesting</i>			Summer rain is erratic. Dig more number of jalkund or pond to store water right now. Harvested rainwater can give you off season vegetable and a good price

### *Horticultural crop*

- Standing rain water in the field may be drained off immediately after rain
- Planting of vegetable seedlings may be done in raised bed to avoid the soil borne diseases / damping
- Soil application of biocontrol agents (Trichoderma) at 2.5kg/ha with farm manure before transplanting
- Transplanting of King chili should be done

<i>Khasi Mandarin</i>		Leaf miner, aphids, mites, trunk	Spray neem oil @5ml/litre water against leaf miner, aphids, mites. Shake the trees to collect adult trunk borer
-----------------------	--	----------------------------------	---

			borer	
	<b>Cucurbits</b>	Vegetative stage		<i>Provide Proper staking</i>
	<b>Okra</b>	Flowering stage	-	<i>Timely earthing up should be done</i>
	<b>Brinjal</b>	Field preparation		<i>Thoroughly prepare the field with application of well dried FYM. Maize should be planted on the border to control movement of white flies</i>
	<b>Ginger / turmeric</b>	Sowing	-	<i>Before sowing, treat ginger rhizome in 5% (5 kg in 100 litre of water) Biozin containing Pseudomonas aeruginosa, Trichoderma harzianum and Bacillus brevis for 1 hr.</i>
<b>Livestock</b>				
<ul style="list-style-type: none"> <li>Water stagnation should be avoided in livestock shed to avoid mosquito breeding</li> </ul>				
	<b>Poultry</b>	<ul style="list-style-type: none"> <li>Double the watering space in poultry houses during summer. Provide plenty of fresh , clean and cool drinking water during hot weather</li> <li>In order to encourage feed intake during summer, reduce the energy content of the feed by 10%</li> <li>Birds must be vaccinated against infectious diseases</li> <li>Remove caked up and wet litter immediately from the poultry house to stop excess heat production</li> </ul>		
	<b>Piggery</b>	<ul style="list-style-type: none"> <li>Ensure that pigs have access to an abundant fresh, cool drinking water to avoid heat stress</li> <li>Offer feed that is higher in protein and vitamins to help compensate for their reduction in food intake as pigs tend to eat less feed during summer</li> </ul>		
<b>Fisheries</b>				
		<ul style="list-style-type: none"> <li>Lime pond as per recommended dose @ 10kg/bigha</li> <li>Stock fish fingerling @ 8000/ha</li> <li>Strengthen embankment to avoid erosion due to heavy rainfall</li> <li>If possible, stock bigger size fish seed to ensure faster growth</li> </ul>		

## SCIENTIFIC EXPERT COMMITTEE

Sl.no	Name	Designation	Department	
1	Dr.D.J. Rajkhowa	Principle Scientist	Agronomy	djrajkhowa@gmail.com
2	Dr. L.K. Baishya	Senior Scientist	Agronomy	lkbicar@gmail.com
3	Ph. Romen Sharma	Scientist	Agricultural Extension	romen.agext@gmail.com
4	Dr. Rajesha G	Scientist	Plant Pathology	rajeshag337@gmail.com
5	Dr. Mahak Singh	Scientist	Animal Reproduction & Gynaecology	mahaksinghivri@gmail.com
6	Dr. Azeze Seyie	Scientist	Spices, Plantation & Medicinal & Aromatic Plants	azezeseyie@yahoo.com
7	Jyotish Barman	Scientist	Fisheries Resource Management	jyotish5@gmail.com
8	Aabon W Yanthan	Scientist	Vegetable Science	aabon.iari@gmail.com

