



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Dimapur	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26^oC to 27^oC and 15^oC to 16^oC, respectively. ▪ Relative humidity varied from 52% to 93%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 26^oC - 27^oC and min temp 13^oC to 15^oC ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 47% to 92%. ▪ Wind speed may reach upto 4-5 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

- Sky will be mainly clear. Farmers are advised for application of manure and fertilizer in orchards
- Marigold should be grown as border crop for chilli and tomato as a trap crop
- Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.
- Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Ripening stage		As the crop will be harvested by the coming week keep the threshing floor ready.
<i>Toria/Mustard</i>	Vegetative stage	-	Daily monitoring should be done for pest and disease infestation
<i>Rabi maize</i>	Vegetative stage		Light hoeing should be done when necessary for better control of weeds
<i>Assam lemon</i>			Weeding should be done frequently. Daily monitoring should be done for aphid
Cabbage	Transplanted	Cabbage butterfly/caterpillar	In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation
Broccoli	Transplanted		In the present weather timely irrigation should be done
Radish, carrot	Transplanted		Timely earthing up should be done after weeding
Poultry	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide protein rich ration and ad libitum clean drinking water		
Fishery	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish		



“GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Kiphire	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26⁰C to 27⁰C and 14⁰C to 16⁰C, respectively. ▪ Relative humidity varied from 30% to 79%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 27⁰C - 28⁰C and min temp 13⁰C to 14⁰C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 23% to 74%. ▪ Wind speed may reach upto 2-4 kmph ▪ Wind may blow from easterly direction <p style="background-color: #ffffcc; margin-top: 5px;">Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Sky will be mainly clear.

Farmers are advised for application of manure and fertilizer in mango and other orchard gardens

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Ripening stage		<i>As the crop will be harvested by the coming week keep the threshing floor ready.</i>
Soybean	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
Rabi maize	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Radish, turnip, Carrot	Transplanted		<i>In the present weather, maintain proper moisture condition in the soil. Never irrigate heavily as it will lead to excessive foliage, produce low quality roots and result in late maturity</i>
Pea, Rajmash	Germination stage		<i>Keeping weather in view, try to keep the plot weed free for proper growth</i>
Tomato	Transplanting stage	Leaf curl	<i>This disease is transmitted by white fly. Severe stunting of plants with downward rolling and crinkling of leaves, older leaves becomes brittle and thick. Try to apply carbofuran @ 1.5 kg /ha at the time of transplanting.</i>
Poultry	As the temperature is decreasing, the shed should be protected from cold wind by hanging curtain around the side of the wall or shed. If possible there should be separate space for different age group of birds. In the present weather, Sick bird should be immediately separated/ culled from healthy flock.		
Piggery	Observe their general stage of health, check for parasites and injuries, and watch out for any abnormal feeding habits		
Fishery	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Kohima	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 25⁰C to 27⁰C and 14⁰C to 13⁰C, respectively. ▪ Relative humidity varied from 51% to 94%. ▪ Wind speed ranged from 2 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 50% to 97%. ▪ Wind speed may reach upto 4-5 kmph ▪ Wind may blow from south easterly direction <p style="background-color: #ffffcc; margin-top: 5px;">Weekly cumulative Rainfall: 0 mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Sowing and transplanting should be carried out
 Marigold should be grown as border crop for chilli and tomato as a trap crop

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Ripening stage		<i>As the crop will be harvested by the coming week keep the threshing floor ready.</i>
<i>Soybean</i>	Harvesting stage		<i>In the present weather, harvesting of the pods should be done. Those that are harvested should be dried immediately during clear day</i>
<i>Toria/Mustard</i>	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Pea</i>	Sowing		<i>Keeping view in weather, sowing of pea should not be delayed, because late sowing will reduces seed yield and will exposes the crop to insect damage.</i>
<i>Khasi mandarin</i>			<i>For good yield , maintain proper sanitation in the orchard</i>
<i>Winter vegetables</i>	Transplanting stage		<i>In the present weather transplanting of vegetables should be carried out.</i>
<i>Poultry</i>	Chickens and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks.		
<i>Piggery</i>	Provide high energy feeds like maize + wheat bran during cold days. Always disinfect the animal shed regularly.		
<i>Fishery</i>	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Longleng	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26⁰C to 27⁰C and 14⁰C to 16⁰C, respectively. ▪ Relative humidity varied from 45% to 80%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 26⁰C - 28⁰C and min temp 13⁰C to 15⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 48% to 89%. ▪ Wind speed may reach upto 2-3 kmph ▪ Wind may blow from south easterly direction

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Marigold should be grown as border crop for chilli and tomato as a trap crop

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done after 7th at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
<i>Soybean</i>	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
<i>Toria/Mustard</i>	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Khasi Mandarin</i>			<i>As flowering has started avoid spraying of insecticides</i>
<i>Rajmash</i>	Vegetative stage		<i>Proper mulching should be carried out</i>
<i>Cabbage</i>	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
<i>Tomato</i>	Transplanting stage		<i>In the present weather as transplanting is going on, maintain recommended spacing for good yield</i>
<i>Poultry</i>			<i>Vaccinate against prevalent disease like Ranikhet disease and deworm every two months for backyard and intensive system both</i>
<i>Piggery</i>			<i>Daily inspection early in the morning should be done for any abnormal behaviour</i>
<i>Fishery</i>			<i>Prune tree branches around the pond so that leaves do not fall on water. Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Mokokchung	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24⁰C to 25⁰C and 14⁰C to 15⁰C, respectively. ▪ Relative humidity varied from 42% to 89%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 49% to 84%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Marigold should be grown as border crop for chilli and tomato as a trap crop

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
Soybean	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
Rabi maize	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Khasi Mandarin			<i>As flowering has started avoid spraying of insecticides</i>
Rajmash	Vegetative stage		<i>Proper mulching should be carried out</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Tomato	Transplanting stage		<i>In the present weather as transplanting is going on, maintain recommended spacing for good yield</i>
Poultry			<i>Vaccinate against prevalent disease like Ranikhet disease and deworm every two months for backyard and intensive system both</i>
Piggery			<i>Daily inspection early in the morning should be done for any abnormal behaviour</i>
Fishery			<i>Prune tree branches around the pond so that leaves do not fall on water. Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Mon	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 25⁰C to 26⁰C and 14⁰C to 15⁰C, respectively. ▪ Relative humidity varied from 42% to 85%. ▪ Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 50% to 92%. ▪ Wind speed may reach upto 3- 4 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES
General Advisories: Sowing and transplanting should be carried out

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
<i>Soybean</i>	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
<i>Potato</i>	Sowing stage		<i>Sowing of potato should be done as soon as possible. Seed treatment with 2 gm Mancozeb and 1 gm Carbendazim mixed per litre of water per kilogram of seed for five minute should be done. Thereafter, seeds should be dried in shade before planting in the field</i>
<i>Toria/Mustard</i>	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
<i>Pea</i>	Sowing		<i>Keeping view in weather, sowing of pea should not be delayed, because late sowing will reduces seed yield and will exposes the crop to insect damage.</i>
<i>Rabi maize</i>	Vegetative stage		<i>Plants should be earthed up after every weeding.</i>
<i>Citrus</i>		Trunk borer	<i>To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in petrol or kerosene and plug with mud.</i>
<i>Vegetables Cabbage, tomato, radish, brocolli</i>	Transplanted		<i>In the present weather Timely irrigation should be done</i>
<i>Rajmash</i>	Vegetative stage		<i>In the present weather condition, Proper mulching should be done</i>
<i>Poultry</i>			<i>Vaccinate against prevalent disease like Ranikhet disease and deworm every two months for backyard and intensive system both</i>
<i>Piggery</i>			<i>Daily inspection early in the morning should be done for any abnormal behaviour</i>
<i>Fishery</i>			<i>If the water level is very low, partially harvest some of the stocked fishes</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Peren	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26⁰C to 27⁰C and 15⁰C to 16⁰C, respectively. ▪ Relative humidity varied from 39% to 88%. ▪ Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 39% to 88%. ▪ Wind speed may reach upto 3- 4 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

- Sky will be mainly clear. Farmers are advised for application of manure and fertilizer in orchards
- Marigold should be grown as border crop for chilli and tomato as a trap crop
- Toria and linseed should be sown as second crop utilizing the residual moisture in the field after the harvest of early to medium duration paddy varieties.
- Sow zero tillage lentil and pea after rice by opening small furrow between rice stubbles

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Ripening stage		As the crop will be harvested by the coming week keep the threshing floor ready.
<i>Toria/Mustard</i>	Vegetative stage	-	Daily monitoring should be done for pest and disease infestation
<i>Rabi maize</i>	Vegetative stage		Light hoeing should be done when necessary for better control of weeds
<i>Assam lemon</i>			Weeding should be done frequently. Daily monitoring should be done for aphid
Cabbage	Transplanted	Cabbage butterfly/caterpillar	In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation
Broccoli	Transplanted		In the present weather timely irrigation should be done
Radish, carrot	Transplanted		Timely earthing up should be done after weeding
Poultry	As the temperature is dropping, provide vitamin complex. Get ready to provide warmth by use of incandescent bulbs to protect them from cold winds by use of plastic sheets, jute sacks and plastic sacks		
Piggery	Protect the animals from cold draft using gunny bags and other locally available materials Provide protein rich ration and ad libitum clean drinking water		
Fishery	If the pond has minimum 1m depth water, apply lime @ 50kg/bigha along with raw cow dung @ 125 kg/bigha, Urea 3.5 kg/bigha, SSP 2.7 kg /bigha. If water depth is less than 1m harvest 50% of the stocked fish		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Phek	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 26⁰C to 28⁰C and 14⁰C to 16⁰C, respectively. ▪ Relative humidity varied from 43% to 87%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 23⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be mainly clear the coming week ▪ Relative Humidity is likely to range from 32% to 87%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from mostly south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Sowing and transplanting should be carried out
 Marigold should be grown as border crop for chilli and tomato as a trap crop

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Ripening stage		<i>As the crop will be harvested by the coming week keep the threshing floor ready.</i>
<i>Soybean</i>	Harvesting stage		<i>In the present weather, harvesting of the pods should be done. Those that are harvested should be dried immediately during clear day</i>
<i>Toria/Mustard</i>	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Pea</i>	Sowing		<i>Keeping view in weather, sowing of pea should not be delayed, because late sowing will reduces seed yield and will exposes the crop to insect damage.</i>
<i>Khasi mandarin</i>			<i>For good yield , maintain proper sanitation in the orchard</i>
<i>Winter vegetables</i>	Transplanting stage		<i>In the present weather transplanting of vegetables should be carried out.</i>
<i>Poultry</i>	Chickens and brooding chicks should be provided warmth by use of incandescent bulbs and protection from cold winds by use of plastic sheets, jute sacks and plastic sacks.		
<i>Piggery</i>	Provide high energy feeds like maize + wheat bran during cold days. Always disinfect the animal shed regularly.		
<i>Fishery</i>	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Tuensang	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 24⁰C to 27⁰C and 14⁰C to 15⁰C, respectively. ▪ Relative humidity varied from 45% to 79%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain the coming week. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 24% to 81%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories: Sky will be mainly clear.

Farmers are advised for application of manure and fertilizer in mango and other orchard gardens

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
TRC/WRC paddy	Ripening stage		<i>As the crop will be harvested by the coming week keep the threshing floor ready.</i>
Soybean	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
Toria/Mustard	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
Rabi maize	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
Cabbage	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
Radish, turnip, Carrot	Transplanted		<i>In the present weather, maintain proper moisture condition in the soil. Never irrigate heavily as it will lead to excessive foliage, produce low quality roots and result in late maturity</i>
Pea, Rajmash	Germination stage		<i>Keeping weather in view, try to keep the plot weed free for proper growth</i>
Tomato	Transplanting stage	Leaf curl	<i>This disease is transmitted by white fly. Severe stunting of plants with downward rolling and crinkling of leaves, older leaves becomes brittle and thick. Try to apply carbofuran @ 1.5 kg /ha at the time of transplanting.</i>
Poultry	As the temperature is decreasing, the shed should be protected from cold wind by hanging curtain around the side of the wall or shed. If possible there should be separate space for different age group of birds. In the present weather, Sick bird should be immediately separated/ culled from healthy flock.		
Piggery	Observe their general stage of health, check for parasites and injuries, and watch out for any abnormal feeding habits		
Fishery	If the water level is very low, partially harvest some of the stocked fishes		



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Wokha	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 25⁰C to 26⁰C and 14⁰C to 15⁰C respectively. ▪ Relative humidity varied from 62% to 89%. ▪ Wind speed ranged from 2 to 3 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 55% to 89%. ▪ Wind speed may reach upto 2- 3 kmph ▪ Wind may blow from easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Marigold should be grown as border crop for chilli and tomato as a trap crop

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
<i>Soybean</i>	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
<i>Toria/Mustard</i>	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Khasi Mandarin</i>			<i>As flowering has started avoid spraying of insecticides</i>
<i>Rajmash</i>	Vegetative stage		<i>Proper mulching should be carried out</i>
<i>Cabbage</i>	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
<i>Tomato</i>	Transplanting stage		<i>In the present weather as transplanting is going on, maintain recommended spacing for good yield</i>
<i>Poultry</i>			<i>Vaccinate against prevalent disease like Ranikhet disease and deworm every two months for backyard and intensive system both</i>
<i>Piggery</i>			<i>Daily inspection early in the morning should be done for any abnormal behaviour</i>
<i>Fishery</i>			<i>Prune tree branches around the pond so that leaves do not fall on water. Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface</i>



GRAMIN KRISHI MAUSAM SEWA
ICAR RESEARCH COMPLEX FOR NEH REGION
Nagaland Center, Medziphema-797106
Dimapur, Nagaland



Telephone No: 03862-247241 (O) / Email: amfu.jharnapani@gov.in

Date: 13.11.18	District: Zunheboto	Bulletin No: 91/2018
Weather of preceding week		Weather forecast valid upto 18.11.2018
<ul style="list-style-type: none"> ▪ No rain occurred the past week ▪ Max and min temperatures ranged 25⁰C to 26⁰C and 13⁰C to 15⁰C, respectively. ▪ Relative humidity varied from 45% to 85%. ▪ Wind speed ranged from 3 to 4 kmph 		<ul style="list-style-type: none"> ▪ Probability of no rain. ▪ Max temp is likely to be 21⁰C - 22⁰C and min temp 10⁰C to 11⁰C ▪ Sky is likely to be clear the coming week ▪ Relative Humidity is likely to range from 45% to 84%. ▪ Wind speed may reach upto 3-4 kmph ▪ Wind may blow from south easterly direction <p>Weekly cumulative Rainfall: 0mm</p>

WEATHER FORECAST BASED AGROMET ADVISORIES

General Advisories:

Marigold should be grown as border crop for chilli and tomato as a trap crop

Main crop	Growth Stage	Pest/Disease	Weather based Agro-Advisories
<i>TRC/WRC paddy</i>	Harvesting stage		<i>Keeping weather in view, the harvesting of mature rice crop should be done at the earliest. After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done</i>
<i>Soybean</i>	Harvesting stage		<i>Suitable time for harvesting of the pods. It should be properly dried before storing</i>
<i>Toria/Mustard</i>	Vegetative stage	-	<i>Daily monitoring should be done for pest and disease infestation</i>
<i>Rabi maize</i>	Vegetative stage		<i>Light hoeing should be done when necessary for better control of weeds</i>
<i>Khasi Mandarin</i>			<i>As flowering has started avoid spraying of insecticides</i>
<i>Rajmash</i>	Vegetative stage		<i>Proper mulching should be carried out</i>
<i>Cabbage</i>	Transplanted	Cabbage butterfly/caterpillar	<i>In the present weather monitoring should be done for cabbage caterpillar. Hand picking and destruction of egg mass and young larvae or apply 4% Neem Seed Kernel Extract (NSKE) and spray 0.05 % Malathion or quinalphos in heavy infestation</i>
<i>Tomato</i>	Transplanting stage		<i>In the present weather as transplanting is going on, maintain recommended spacing for good yield</i>
<i>Poultry</i>			<i>Vaccinate against prevalent disease like Ranikhet disease and deworm every two months for backyard and intensive system both</i>
<i>Piggery</i>			<i>Daily inspection early in the morning should be done for any abnormal behaviour</i>
<i>Fishery</i>			<i>Prune tree branches around the pond so that leaves do not fall on water. Do take care of your pond every day when you take food to your fish, make sure the pond remains full of water, do not let weeds cover more than one quarter of the surface</i>