



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of DIMAPUR(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	25	12	80	38	5.4	109	0
2020-02-13	0	27	12	74	36	5.2	109	0
2020-02-14	0	28	13	69	34	5.2	22	0
2020-02-15	0	29	14	62	32	5.3	124	0
2020-02-16	0	29	13	65	29	5.5	109	0

### Weather Summary/Alert:

Probability of no rainfall the coming week
Etia lakha somoi dhe pani nigheribo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato
Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement
Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : Rice should be stored in paddy form rather than milled rice as the husk provides some protection against insects
	STORAGE : Dhan tho sob nukhundighenah, kiman lageh eman hi khundibi. Sob tho khundighenah rakhithakileh puka bi lagijabo
GINGER	LAND PREPARATION : Start preparation of land
	LAND PREPARATION : Mathi bonaboleh suru kuribi

Crop(Varieties)	Crop Specific Advisory
OKRA/ LADIES' FINGER	LAND PREPARATION : Start land preparation. Well decomposed FYM or organic manure or poultry manure can be incorporated at the time of land preparation
	LAND PREPARATION : Mathi bonaboleh suru kuribi. Mathi bona somoi tae bhal parah sukhijah FYM, organic manure or murgi manure milabo pariboh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
VEGETABLE PEA	GERMINATING STAGE: In the present weather as there will be no rain, light irrigation should be provided
	GERMINATING STAGE: Etia lakha somoi dhe pani nigheribo dho, pani somoi dhe alibi
BRINJAL	NURSERY: Start nursery preparation. To avoid mortality of seedlings due to damping off, drenching of the beds with Bavistin (15-20 g/10 litres of water) is effective.
	NURSERY: Nursery bonaboleh suru kuribi. Bijon guti puchighenah murinajabo karoneh Bavistin (15-20 g/10 litres pani)
LEMON	Timely remove the dry twigs and dry and diseased leaves
	Somoi tae sukha branch, sukha aru bimar pata khan hatabi

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW	Adequate measures should be taken to protect the animals from the wet weather, as well as the drop in temperature when the skies clear after the rain.
	Din biyah para janwar khan bachaighenah rakhibi
PIG	Monitored for signs of discomfort (extensive shivering, weakness, lethargy, etc.). Special attention should be paid to very young and old animals because they may be less able to tolerate temperature extremes
	Gahori khan komjori aru kapithaka sabo lageh. Bacha aru omor bishi gahori khan bhal parah sabi. Thanda gorom bishi hua tho lobo napareh

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	If complete removal of water is not possible, remove the aquatic vegetation manually. Go for repeated netting to remove all small and unwanted fishes.
	Pani sob phekaboleh naparilehbi, jongol ghas khan sapha kuribi aru jal marighenah chutu aru nalagia mass khan sob hatabi

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
FOWL	For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.
	Bimar nalagibo karoneh sapha pani thik thak dibi, achanak parah dana bodli nukuribi, ghor tae hawa aha jah thakibo lageh, ekta jagah tae murgi jama kurighenah narikhibi aru, murijah khan jolaidibi nahoileh mati tae rupaidibi



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of KIPHIRE(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	26	10	77	21	3.3	68	0
2020-02-13	0	28	12	68	21	3.0	64	0
2020-02-14	0	30	12	68	18	3.1	64	0
2020-02-15	0	30	12	74	24	3.5	45	0
2020-02-16	0	27	11	64	24	3.8	64	0

### Weather Summary/Alert:

Probability of no rainfall the coming days

Pani etia ahibo lhe hapta dho nigheri bo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Use rodent traps and other means to control rodent, insect and pests
	STORAGE: Chuha aru puka komti kuriboleh chuha phang aru alag opai cholabi
MAIZE	LAND PREPARATION : Plan for maximizing plant diversity by inter cropping of maize with suitable pulse crops of your particular region. Eg: Maize + pigeon pea/black gram /green gram. Plant Napier grass in the border rows to act as FAW trap crop.

Crop(Varieties)	Crop Specific Advisory
	LAND PREPARATION : Nijor jagah hisap tae makoi logotae alag pulse sobji lagabi. Eg: Makoi logotae pigeon pea/black gram /green gram. FAW trap crop karoneh kinar khan tae Napier grass lagabi

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	SOWING: Sowing should be started. Before sowing, well decomposed FYM should be mixed with soil during last ploughing
	SOWING: Bijon lagaboleh suru kuribi. Nalagah agotae last khunda somoi tae mathi tae bhal parah sukija FYM milabi
CABBAGE	HARVESTING STAGE: Harvesting should not be delayed as even a few days beyond maturity can result in split heads and increased incidence of field disease.
	HARVESTING STAGE: Kobi thik somoi tae harvest kurilobi. Diri kurileh pata khan khuli jabo aru bimar bi lagijobo parah.
BANANA	Keep the soil loose by harrowing from time to time
	Somoi tae kol ghas kinar mathi khan dhilah kurighenah rakhibi

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a
	Sapha parah gahori palia parah gahori gao bhal thake. Latrine hudai dapha kurah, ghor sapha rakha.

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Harvest the previously stocked fish preferably by 1st week
	Puilah tae dhalia mass khan etu hopta tae dhurilobi

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Poultry house should be rain-proof and protected against predators. It should have windows with wire mesh for adequate ventilation. Dusty environment irritates the respiratory tract of the chicks. Besides dust is one of the vehicles of transmission of diseases
	Murgi ghor tae borokhon nahibo hisap tae aru murgi ke khaboleh janwar khan ahiboleh naparah hisap tae bonabi. Hawa bhal parah ahiboleh iron jali lagabi. Ghor tae dhul nahibo jagah tae bonabi. Dhul parah murigi sas lua duk pai



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of KOHIMA(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	25	11	98	53	4.9	162	1
2020-02-13	0	26	12	97	47	5.0	162	0
2020-02-14	0	28	13	88	41	5.1	292	0
2020-02-15	0	29	12	84	43	6.1	206	0
2020-02-16	0	27	13	82	44	6.0	158	0

### Weather Summary/Alert:

Probability of no rainfall the coming days

Pani etia ahibolhe hapta dho nigheri bo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement.

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : If there are cracks or holes in the storage room then sealed it, to avoid entry of water by seepage from ground or walls Pani khan ghor bitor tae nahibo karoneh chenda khan bon kuribi
MAIZE	LAND PREPARATION: Start field preparation . The soil should be well drained . Makhoi lagabo lhe mati bonabo shoru khoribi. Mati dho pani pal bara jabo lhe laghe dhe hei lagabi
GINGER	LAND PREPARATION : Land preparation should be started

Crop(Varieties)	Crop Specific Advisory
	LAND PREPARATION : mathi bonaboleh suru kuribi

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	Plantation should be protected from strong winds by growing tall plants along the farm border. Hawa jor parah kol ghas biya nukuriboleh bajan kinar tae lamba ghas lagabi
POTATO	LAND PREPARATION : Start preparation of the main field. Only certified seed free from diseases should be sown. LAND PREPARATION : Mathi bonaboleh suru kuribi. Bijon dho certified hobo lakhe aru bimar natakha hei jolabo laghe
CABBAGE	NURSERY PREPARATION : Beds should be raised to avoid water logging in heavy soils NURSERY PREPARATION : Pani jama nahobo karoneh mati uthabi

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Piglets should be given iron injection on the 4th and 14th day of farrowing, deep intramuscularly to prevent piglet anaemia. Oral iron supplement may be given Gahori bacha khan jonom hughenah 4 aru 14 din tae iron injection maribi. Muk tae khela iron suppliment bi dibo parah

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Remove all small and unwanted fishes from the pond as they compete for food, space and oxygen with the stocked fishes. Chutu aru nalagia mass khan pukhuri parah hatabi. Etu mass khan dana , jagah aru oxygen karoneh alagh mass logotae jujithakeh

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BREEDER	Vanaraja, a dual purpose bird is suitable for backyard poultry farming due to its high production potential in terms of meat and egg than that of local/desi bird (low produces of meat and egg). It thrives well under low feeding and managerial practices due to its adaptability and local climatic conditions Vanaraja murgi khan local/desi murgi khan parah mungso tae bishi hoi aru anda bi bishi de. Dana komti dilibi aru bhal parah nasailebi, amikhan lagah jagah tae bhal hoi



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

**Weather Forecast of LONGLENG(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)**

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0.0	16.0	7.0	80	37	5.0	158	0
2020-02-13	0.0	17.0	7.0	72	31	5.1	154	0
2020-02-14	0.0	17.0	7.0	63	25	5.1	154	0
2020-02-15	0.0	17.0	7.0	68	30	5.0	158	1
2020-02-16	0.0	18.0	8.0	71	34	5.4	135	0

### Weather Summary/Alert:

Probability of no rainfall the coming days. Sky will be clear

Etia ahiboleh din dho pani nikiriboh leh pareh hoilebi

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement.

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : The godown where the grains has been kept should be clean, dry and properly aerated
	Dhan rakha ghor sukha aru sapha hobo lageh aru hawah aha jah thakibo lageh
MAIZE	LAND PREPARATION :One deep ploughing should be given, followed by two or three harrowings to bring the soil to a fine tilth
	LAND PREPARATION :Mathi bhal parah khundibi

**Horticulture Specific Advisory:**

<b>Horticulture (Varieties)</b>	<b>Horticulture Specific Advisory</b>
COLOCASIA/ TARO	LAND PREPARATION : Start preparing the land. Plough the field to a fine tilth and form ridges and furrows
	LAND PREPARATION : Mathi khundiboleh suru kuribi aru bhal parah khundibi
CHILLI	NURSERY PREPARATION : Treat the seeds with Trichoderma viride @ 4 g / kg or Pseudomonas fluorescens @ 10 g/ kg and sow in lines spaced at 10 cm in raised nursery beds and cover with sand. Watering with rose can has to be done daily.
	NURSERY PREPARATION : Bijon khan Trichoderma viride @ 4 g / kg nahoileh Pseudomonas fluorescens @ 10 g/ kg parah traet kurilobi aru 10 cm raised nursery beds lines spaced tae lagabi aru mathi parah bon kuribi. Hudai rose can parah pani dibi
MANDARIN ORANGE	POST HARVEST : Post-harvest handling of fruits must be followed carefully to avoid any sort of injury of fruits.
	POST HARVEST : Fruits khan harvest kurah pasotae biyah nahoboleh bhal parah chubi
TAPIOCA	LAND PREPARATION: Well drained soil should be selected
	LAND PREPARATION: Bhal parah pani jah lagha mathi hea dhe lagabi
CABBAGE	HARVESTING STAGE: Harvested produce should always be stored in shade before packing.
	Harvest kurighenah packing nukurah agotae, thanda jagah tae rakhibi

**Livestock Specific Advisory:**

<b>Livestock(Varieties)</b>	<b>Livestock Specific Advisory</b>
PIG	Keep floor dry to avoid skin disease
	Skin bimar nulabo karoneh ghor lagah ghor sukha rakhibi

**Fisheries Specific Advisory:**

<b>Fisheries(Varieties)</b>	<b>Fisheries Specific Advisory</b>
COMMON FISH	Complete dewatering of pond is recommended. It shall ensure complete removal of fish and unwanted vegetation. After dewatering remove the pond bottom mud and the excavated mud can be used for strengthening the pond dyke. Expose the pond bottom to bright sun for about 15-20 days
	Pukhuri parah pani sob phekailebi. Nalagah mass aru ghas khan pukuri tae thakibo nidibi. Pani phekaighenah nichea lagah mathi hatabi aru etu mathi parah pukhuri lagah opor kinar khan bonabi. 15-20 din karoneh pukhuri tho dhup lagibo dibi

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
FOWL	Poultry house should have windows with wire mesh for adequate ventilation. Dusty environment irritates the respiratory tract of the chicks. Besides dust is one of the vehicles of transmission of diseases.
	Hawa bhal parah ahiboleh iron jali lagabi. Ghor tae dhul nahibo jagah tae bonabi. Dhul parah murigi sas lua duk pai.



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of MOKOKCHUNG(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	25	11	84	41	4.6	135	0
2020-02-13	0	27	12	79	38	4.4	153	0
2020-02-14	0	28	12	71	32	4.6	154	0
2020-02-15	0	28	14	74	33	4.6	154	1
2020-02-16	0	27	13	71	35	5.0	112	0

### Weather Summary/Alert:

Probability of no rainfall the coming days. Sky will be clear

Etia ahibleh din dho pani nikiriboh leh parah hoilebi

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : The godown where the grains has been kept should be clean, dry and properly aerated
	Dhan rakha ghor sukha aru sapha hobo lageh aru hawah aha jah thakibo lageh
MAIZE	LAND PREPARATION : One deep ploughing should be given, followed by two or three harrowings to bring the soil to a fine tilth
	LAND PREPARATION : Mathi bhal parah khundibi

**Horticulture Specific Advisory:**

Horticulture (Varieties)	Horticulture Specific Advisory
COLOCASIA/ TARO	LAND PREPARATION : Start preparing the land. Plough the field to a fine tilth and form ridges and furrows
	LAND PREPARATION : Mathi khundiboleh suru kuribi aru bhal parah khundibi
CHILLI	NURSERY PREPARATION : Treat the seeds with Trichoderma viride @ 4 g / kg or Pseudomonas fluorescens @ 10 g/ kg and sow in lines spaced at 10 cm in raised nursery beds and cover with sand. Watering with rose can has to be done daily.
	NURSERY PREPARATION : Bijon khan Trichoderma viride @ 4 g / kg nahoileh Pseudomonas fluorescens @ 10 g/ kg parah traet kurilobi aru 10 cm raised nursery beds lines spaced tae lagabi aru mathi parah bon kuribi. Hudai rose can parah pani dibi
MANDARIN ORANGE	POST HARVEST : Post-harvest handling of fruits must be followed carefully to avoid any sort of injury of fruits.
	POST HARVEST : Fruits khan harvest kurah pasotae biyah nahoboleh bhal parah chubi
TAPIOCA	LAND PREPARATION: Well drained soil should be selected
	LAND PREPARATION: Bhal parah pani jah lagha mathi hea dhe lagabi
CABBAGE	HARVESTING STAGE: Harvested produce should always be stored in shade before packing
	Harvest kurighenah packing nukurah agotae, thanda jagah tae rakhibi

**Livestock Specific Advisory:**

Livestock(Varieties)	Livestock Specific Advisory
PIG	Keep floor dry to avoid skin disease
	Skin bimar nulabo karoneh ghor lagah ghor sukha rakhibi

**Fisheries Specific Advisory:**

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Complete dewatering of pond is recommended. It shall ensure complete removal of fish and unwanted vegetation. After dewatering remove the pond bottom mud and the excavated mud can be used for strengthening the pond dyke. Expose the pond bottom to bright sun for about 15-20 days
	Pukhuri parah pani sob phekailobi. Nalagah mass aru ghas khan pukuri tae thakibo nidibi. Pani phekaighenah nichea lagah mathi hatabi aru etu mathi parah pukhuri lagah opor kinar khan bonabi. 15-20 din karoneh pukhuri tho dhup lagibo dibi

**Poultry Specific Advisory:**

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Poultry house should have windows with wire mesh for adequate ventilation. Dusty environment irritates the respiratory tract of the chicks. Besides dust is one of the vehicles of transmission of diseases.
	Hawa bhal parah ahiboleh iron jali lagabi. Ghor tae dhul nahibo jagah tae bonabi. Dhul parah murigi sas lua duk pai.



## Gramin Krishi Mausam Sewa

India Meteorological Department

ICAR Complex Branch

Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of MON(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	26	11	77	41	5.1	154	0
2020-02-13	0	28	12	69	31	5.4	154	0
2020-02-14	0	28	12	62	25	5.3	158	0
2020-02-15	0	28	13	65	32	5.2	158	1
2020-02-16	0	27	13	70	35	5.3	154	0

### Weather Summary/Alert:

Probability of no rainfall the coming days. Sky will be clear

Etia ahibleh din dho pani nikiriboh leh pareh hoilebi, akash bi sabha hobo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Monitor timely for storage pest like moth and weevil infestation
	STORAGE: Dhah ghor tae pukha moth aru weevil ase nah nai somoi tae sai thakibi
FOXTAIL MILLET	LAND PREPARATION : Start land preparation for sowing
	Mati bonabolhe shuro khoribi
MAIZE	LAND PREPARATION: Deep summer ploughing should be done as it helps in exposing resting stages of pests. To protect seeds from soil borne diseases and insect pest, seed treatment is necessary

Crop(Varieties)	Crop Specific Advisory
	LAND PREPARATION: Mathi phuka khan dhup aparah muriboleh, mathi khunda somoi tae olop dhada khundibi. Bijon guti khan mathi lagah bimar parah bachiboleh seed treatment kuribolageh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
ORANGE	To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in the mixture of kerosene and water(1:1) and plug with mud
	Ghas lakha korithe pukha khan daka tho, wire parah sabha khoribi etu biche de cotton kerosene aru pani mixture (1:1) DHE BICHABI ETU BICHE THE jenta the kusaidhebi. Mati para jenta tho pon koribi
ONION	STORAGE : The storage rots could be checked if proper cleanliness is maintained in store
	Biyas rakha jakha dho sabha rakhibo laghe dai nobojibo karoneh

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Vaccinate all newly born piglets against swine fever at the age of 3 months followed by annual booster dose.
	Gauri baccha din mohinah huwa bicche dhe tho swine fever karoneh vaccine khoriboh laghe

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Remove all small and unwanted fishes from the pond as they compete for food, space and oxygen with the stocked fishes
	Chotu aru noloboh mass khan dhorilobi khelhe khoileh etu khan para bal mass khan khaluwa kharoneh chuche

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	Maintain high hygienic and sanitary measures in poultry by good management to avoid early exposure of young chicken to diseases
	Maintain high hygienic and sanitary measures in poultry by good management to avoid early exposure of young chicken to diseases



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of PEREN(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	25	10	93	39	4.5	112	0
2020-02-13	0	27	12	88	34	4.4	109	0
2020-02-14	0	28	13	69	28	4.3	116	0
2020-02-15	0	29	13	70	33	4.9	158	0
2020-02-16	0	28	13	73	34	4.5	116	0

### Weather Summary/Alert:

Probability of no rainfall the coming days

Pani etia ahibo lhe hapta dho nigheri bo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : Rice should be stored in paddy form rather than milled rice as the husk provides some protection against insects
	STORAGE : Dhan tho sob nukhundighenah, kiman lageh eman hi khundibi. Sob tho khundighenah rakhithakileh puka bi lagijabo
GINGER	LAND PREPARATION : Start preparation of land
	LAND PREPARATION : Mathi bonaboleh suru kuribi

Crop(Varieties)	Crop Specific Advisory
OKRA/ LADIES' FINGER	LAND PREPARATION : Start land preparation. Well decomposed FYM or organic manure or poultry manure can be incorporated at the time of land preparation.
	LAND PREPARATION : Mathi bonaboleh suru kuribi. Mathi bona somoi tae bhal parah sukhijah FYM, organic manure or murgi manure milabo pariboh

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
VEGETABLE PEA	GERMINATING STAGE: In the present weather as there will be no rain, light irrigation should be provided
	GERMINATING STAGE: Etia lakha somoi dhe pani nigheribo dho, pani somoi dhe alibi
BRINJAL	NURSERY: Nursery bonaboleh suru kuribi. Bijon guti puchighenah murinajabo karoneh Bavistin (15-20 g/10 litres pani)
	NURSERY: Nursery bonaboleh suru kuribi. Bijon guti puchighenah murinajabo karoneh Bavistin (15-20 g/10 litres pani)
LEMON	Timely remove the dry twigs and dry and diseased leaves
	Somoi tae sukha branch, sukha aru bimar pata khan hatabi

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW	Adequate measures should be taken to protect the animals from the wet weather, as well as the drop in temperature when the skies clear after the rain.
	Din biyah para janwar khan bachaighenah rakhibi
PIG	Monitored for signs of discomfort (extensive shivering, weakness, lethargy, etc.). Special attention should be paid to very young and old animals because they may be less able to tolerate temperature extremes
	Gahori khan komjori aru kapithaka sabo lageh. Bacha aru omor bishi gahori khan bhal parah sabi. Thanda gorom bishi hua tho lobo napareh

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	If complete removal of water is not possible, remove the aquatic vegetation manually. Go for repeated netting to remove all small and unwanted fishes.
	Pani sob phekaboleh neparilehbi, jongol ghas khan sapha kuribi aru jal marighenah chutu aru nalagia mass khan sob hatabi

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
FOWL	For prevention of diseases, some general measures should be followed. Provide ample quantity of clean drinking water, don't make sudden changes in feed, maintain good ventilation avoid overcrowding and dispose dead birds by burning or burial.
	Bimar nalagibo karoneh sapha pani thik thak dibi, achanak parah dana bodli nukuribi, ghor tae hawa aha jah thakibo lageh, ekta jagah tae murgi jama kurighenah narikhibi aru, murijah khan jolaidibi nahoileh mati tae rupaidibi



## Gramin Krishi Mausam Sewa

India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



# Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of PHEK(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	26	10	91	32	2.0	240	0
2020-02-13	0	28	12	85	29	2.0	248	0
2020-02-14	0	29	13	80	26	2.4	249	0
2020-02-15	0	30	13	79	30	3.5	257	0
2020-02-16	0	27	12	73	31	3.0	158	0

### Weather Summary/Alert:

Probability of no rainfall the coming days
Pani etia ahibolhe hapta dho nigheri bo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.
Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement
Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	SSTORAGE : If there are cracks or holes in the storage room then sealed it, to avoid entry of water by seepage from ground or walls
	Pani khan ghor bitor tae nahibo karoneh chenda khan bon kuribi
GINGER	LAND PREPARATION : Land preparation should be started
	LAND PREPARATION : Mathi bonaboleh suru kuribi
MAIZE	LAND PREPARATION: Start field preparation . The soil should be well drained

Crop(Varieties)	Crop Specific Advisory
	Makhoi lagabo lhe mati bonabo shoru khoribi. Mati dho pani pal bara jabo lhe laghe dhe hei lagabi

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	Plantation should be protected from strong winds by growing tall plants along the farm border Hawa jor parah kol ghas biya nukuriboleh bajan kinar tae lamba ghas lagabi
POTATO	LAND PREPARATION : Start preparation of the main field. Only certified seed free from diseases should be sown. LAND PREPARATION : Mathi bonaboleh suru kuribi. Bijon dho certified hobo lakhe aru bimar natakha hei jolabo laghe
CABBAGE	NURSERY PREPARATION : Beds should be raised to avoid water logging in heavy soils NURSERY PREPARATION : Pani jama nahobo karoneh mati uthabi

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Piglets should be given iron injection on the 4th and 14th day of farrowing, deep intramuscularly to prevent piglet anaemia. Oral iron supplement may be given Gahori bacha khan jonom hughenah 4 aru 14 din tae iron injection maribi. Muk tae khela iron suppliment bi dibo parah

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Remove all small and unwanted fishes from the pond as they compete for food, space and oxygen with the stocked fishes. Chutu aru nalagia mass khan pukhuri parah hatabi. Etu mass khan dana , jagah aru oxygen karoneh alagh mass logotae jujithakeh

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
BREEDER	Vanaraja, a dual purpose bird is suitable for backyard poultry farming due to its high production potential in terms of meat and egg than that of local/desi bird (low produces of meat and egg). It thrives well under low feeding and managerial practices due to its adaptability and local climatic conditions Vanaraja murgi khan local/desi murgi khan parah mungso tae bishi hoi aru anda bi bishi de. Dana komti dilibi aru bhal parah nasailebi, amikhan lagah jagah tae bhal hoi.



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of TUENSANG(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	25	11	87	32	4.2	112	0
2020-02-13	0	26	11	80	30	4.3	112	0
2020-02-14	0	27	12	70	23	4.2	109	0
2020-02-15	0	28	13	78	28	3.7	112	0
2020-02-16	0	26	12	75	32	4.3	116	0

### Weather Summary/Alert:

Probability of no rainfall the coming days
Pani etia ahibo lhe hapta dho nigheri bo

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.
Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement.
Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Use rodent traps and other means to control rodent, insect and pests
	STORAGE: Chuha aru puka komti kuriboleh chuha phang aru alag opai cholabi
MAIZE	LAND PREPARATION : Plan for maximizing plant diversity by inter cropping of maize with suitable pulse crops of your particular region. Eg: Maize + pigeon pea/black gram /green gram. Plant Napier grass in the border rows to act as FAW trap crop.

Crop(Varieties)	Crop Specific Advisory
	LAND PREPARATION : Nijor jagah hisap tae makoi logotae alag pulse sobji lagabi. Eg: Makoi logotae pigeon pea/black gram /green gram. FAW trap crop karoneh kinar khan tae Napier grass lagabi

### Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	SOWING: Sowing should be started. Before sowing, well decomposed FYM should be mixed with soil during last ploughing
	SOWING: Bijon lagaboleh suru kuribi. Nalagah agotae last khunda somoi tae mathi tae bhal parah sukija FYM milabi
CABBAGE	HARVESTING STAGE: Harvesting should not be delayed as even a few days beyond maturity can result in split heads and increased incidence of field disease.
	HARVESTING STAGE: Kobi thik somoi tae harvest kurilobi. Diri kurileh pata khan khuli jabo aru bimar bi lagijobo parah.
BANANA	Keep the soil loose by harrowing from time to time
	Somoi somoi tae kol ghas kinar mathi khan dhilah kurighenah rakhibi

### Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd
	Sapha parah gahori palia parah gahori gao bhal thake. Latrine hudai dapha kurah, ghor sapha rakha.

### Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Harvest the previously stocked fish preferably by 1st week
	Puilah tae dhalia mass khan etu hopta tae dhurilobi

### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Poultry house should be rain-proof and protected against predators. It should have windows with wire mesh for adequate ventilation. Dusty environment irritates the respiratory tract of the chicks. Besides dust is one of the vehicles of transmission of diseases
	Murgi ghor tae borokhon nahibo hisap tae aru murgi ke khaboleh janwar khan ahiboleh naparah hisap tae bonabi. Hawa bhal parah ahiboleh iron jali lagabi. Ghor tae dhul nahibo jagah tae bonabi. Dhul parah murigi sas lua duk pai



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of WOKHO(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0.0	16.0	7.0	86	45	4.6	109	0
2020-02-13	0.0	17.0	7.0	85	42	4.6	105	0
2020-02-14	0.0	17.0	7.0	78	37	4.6	112	0
2020-02-15	0.0	17.0	7.0	75	36	4.8	110	1
2020-02-16	0.0	18.0	8.0	69	33	4.7	112	0

### Weather Summary/Alert:

Probability of no rainfall the coming days. Sky will be clear

Etia ahiboleh din dho pani nikiriboh leh parah hoilebi,

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : The godown where the grains has been kept should be clean, dry and properly aerated
	Dhan rakha ghor sukha aru sapha hobo lageh aru hawah aha jah thakibo lageh
MAIZE	LAND PREPARATION : One deep ploughing should be given, followed by two or three harrowings to bring the soil to a fine tilth
	LAND PREPARATION : Mathi bhal parah khundibi

**Horticulture Specific Advisory:**

Horticulture (Varieties)	Horticulture Specific Advisory
COLOCASIA/ TARO	LAND PREPARATION :Mathi bhal parah khundibi
	LAND PREPARATION : Mathi khundiboleh suru kuribi aru bhal parah khundibi
CHILLI	NURSERY PREPARATION : Treat the seeds with Trichoderma viride @ 4 g / kg or Pseudomonas fluorescens @ 10 g/ kg and sow in lines spaced at 10 cm in raised nursery beds and cover with sand. Watering with rose can has to be done daily.
	NURSERY PREPARATION : Bijon khan Trichoderma viride @ 4 g / kg nahoileh Pseudomonas fluorescens @ 10 g/ kg parah traet kurilobi aru 10 cm raised nursery beds lines spaced tae lagabi aru mathi parah bon kuribi. Hudai rose can parah pani dibi
MANDARIN ORANGE	POST HARVEST : Post-harvest handling of fruits must be followed carefully to avoid any sort of injury of fruits
	POST HARVEST : Fruits khan harvest kurah pasotae biyah nahoboleh bhal parah chubi
TAPIOCA	LAND PREPARATION: Well drained soil should be selected
	LAND PREPARATION: Bhal parah pani jah lagha mathi hea dhe lagabi
CABBAGE	HARVESTING STAGE: Harvested produce should always be stored in shade before packing.
	Harvest kurighenah packing nukurah agotae, thanda jagah tae rakhibi

**Livestock Specific Advisory:**

Livestock(Varieties)	Livestock Specific Advisory
PIG	Keep floor dry to avoid skin disease
	Skin bimar nulabo karoneh ghor lagah ghor sukha rakhibi

**Fisheries Specific Advisory:**

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Complete dewatering of pond is recommended. It shall ensure complete removal of fish and unwanted vegetation. After dewatering remove the pond bottom mud and the excavated mud can be used for strengthening the pond dyke. Expose the pond bottom to bright sun for about 15-20 days
	Pukhuri parah pani sob phekailobi. Nalagah mass aru ghas khan pukuri tae thakibo nidibi. Pani phekaighenah nichea lagah mathi hatabi aru etu mathi parah pukhuri lagah opor kinar khan bonabi. 15-20 din karoneh pukhuri tho dhup lagibo dibi

**Poultry Specific Advisory:**

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Poultry house should have windows with wire mesh for adequate ventilation. Dusty environment irritates the respiratory tract of the chicks. Besides dust is one of the vehicles of transmission of diseases
	Hawa bhal parah ahiboleh iron jali lagabi. Ghor tae dhul nahibo jagah tae bonabi. Dhul parah murigi sas lua duk pai.



**Gramin Krishi Mausam Sewa**  
India Meteorological Department  
ICAR Complex Branch  
Umiam, Meghalaya



## Agromet Advisory Bulletin

Date : 11-02-2020

Weather Forecast of ZUNHEBOTO(Nagaland) Issued On : 2020-02-11(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-12	0	25	11	95	38	3.9	108	0
2020-02-13	0	27	11	91	36	3.6	108	0
2020-02-14	0	28	12	84	32	3.6	108	0
2020-02-15	0	29	13	88	34	3.5	297	0
2020-02-16	0	26	12	77	34	4.2	112	0

### Weather Summary/Alert:

Probability of no rainfall the coming days. Sky will be clear

Etia ahiboleh din dho pani nikiriboh leh pareh hoilebi

### General Advisory:

Farmers are advised to install pheromone traps @ 3-4 traps per acre for monitoring of the pest population for diamond back moth in cole crop, pod borer in pea and fruit borer in tomato.

Diamond black moth cole crop dhe, pod borer matar dhe aru fruit borer tomato dhe komti kuriboleh, pheromone traps ekta acre tae 3-4 rakhibi

### SMS Advisory:

Keeping increasing temperature in view, farmers are advised for light irrigation in standing crops and vegetables as per requirement

Etia somoi, hawa gorom aru pani nighera somoi dhe, apuni khan bakan dhe pani dibo lakhe

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : The godown where the grains has been kept should be clean, dry and properly aerated
	Dhan rakha ghor sukha aru sapha hobo lageh aru hawah aha jah thakibo lageh
MAIZE	LAND PREPARATION : One deep ploughing should be given, followed by two or three harrowings to bring the soil to a fine tilth
	LAND PREPARATION : Mathi bhal parah khundibi

**Horticulture Specific Advisory:**

Horticulture (Varieties)	Horticulture Specific Advisory
COLOCASIA/ TARO	LAND PREPARATION : Start preparing the land. Plough the field to a fine tilth and form ridges and furrows
	LAND PREPARATION : Mathi khundiboleh suru kuribi aru bhal parah khundibi
CHILLI	NURSERY PREPARATION : Treat the seeds with Trichoderma viride @ 4 g / kg or Pseudomonas fluorescens @ 10 g/ kg and sow in lines spaced at 10 cm in raised nursery beds and cover with sand. Watering with rose can has to be done daily
	NURSERY PREPARATION : Bijon khan Trichoderma viride @ 4 g / kg nahoileh Pseudomonas fluorescens @ 10 g/ kg parah traet kurilobi aru 10 cm raised nursery beds lines spaced tae lagabi aru mathi parah bon kuribi. Hudai rose can parah pani dibi
MANDARIN ORANGE	POST HARVEST : Post-harvest handling of fruits must be followed carefully to avoid any sort of injury of fruits.
	POST HARVEST : Fruits khan harvest kurah pasotae biyah nahoboleh bhal parah chubi
TAPIOCA	LAND PREPARATION: Well drained soil should be selected
	LAND PREPARATION: Bhal parah pani jah lagha mathi hea dhe lagabi
CABBAGE	HARVESTING STAGE: Harvested produce should always be stored in shade before packing.
	Harvest kurighenah packing nukurah agotae, thanda jagah tae rakhibi

**Livestock Specific Advisory:**

Livestock(Varieties)	Livestock Specific Advisory
PIG	Keep floor dry to avoid skin disease
	Skin bimar nulabo karoneh ghor lagah ghor sukha rakhibi

**Fisheries Specific Advisory:**

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Complete dewatering of pond is recommended. It shall ensure complete removal of fish and unwanted vegetation. After dewatering remove the pond bottom mud and the excavated mud can be used for strengthening the pond dyke. Expose the pond bottom to bright sun for about 15-20 days
	Pukhuri parah pani sob phekailobi. Nalagah mass aru ghas khan pukuri tae thakibo nidibi. Pani phekaighenah nichea lagah mathi hatabi aru etu mathi parah pukhuri lagah opor kinar khan bonabi. 15-20 din karoneh pukhuri tho dhup lagibo dibi

**Poultry Specific Advisory:**

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Poultry house should have windows with wire mesh for adequate ventilation. Dusty environment irritates the respiratory tract of the chicks. Besides dust is one of the vehicles of transmission of diseases.
	Hawa bhal parah ahiboleh iron jali lagabi. Ghor tae dhul nahibo jagah tae bonabi. Dhul parah murigi sas lua duk pai.