



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of ZUNHEBOTO(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	2	22	10	98	63	3.2	108	1
2020-02-02	1	24	10	98	43	3.8	112	0
2020-02-03	0	22	8	79	36	3.9	90	0
2020-02-04	0	24	9	97	36	3.4	297	0
2020-02-05	0	24	10	84	32	3.7	112	1

Weather Summary/Alert:

Probability of light rainfall the coming days. Sky will be mainly clear.

Etia ahiboleh din tho pani olop kiribolhe dikaiase

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker

Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh.

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed

Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh
	STORAGE: Pukha khan parah bachibo karoneh, dhan rakha somoi tae neem pata logotae rakhibi

Crop(Varieties)	Crop Specific Advisory
RAPESEED	THRESHING: Cleaned seed must be dried in the sun for four to five days or till the moisture content comes down to 8 percent.
	THRESHING: Bijon guti khan moisture 8% tae nalamah thok dhup tae sukhabhi
TOMATO	Clean and sanitize harvest containers or bins at least weekly, more often if needed, to remove sand, grit, dirt and other residue
	Tomato harvest kura daba khan letra bhalu nathakiboleh, somoi tae sapha kuri thakibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	HARVESTING STAGE: 2 TO 3 pickings should be done at 4-5 days interval
	HARVESTING STAGE: 2 bar parah 3 bar nishinah shingilobi aru shinga majotae 4-5 din charighenah hi shingibi
MANDARIN ORANGE	Harvested fruits should be kept under proper storage condition to ensure that there is no change in the quality even after months of storage
	Harvest kurighenah olop mohinah thok biya habobo karoneh bhal parah rakhibi
CABBAGE	HARVESTING STAGE: Cabbage should be harvested promptly when the heads are firm and mature
	HARVESTING STAGE: Kobi lagah matha dangor huighenah taan hua somoi tae harvest kurilobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Vaccinate all newly born piglets against swine fever at the age of 3 months followed by annual booster dose
	Gahor bacha khan 3 months tae swine fever karoneh vaccinate kurilobi aru etu pichetae bi mari thakibi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Check for any fish disease break out or fish mortality during winter months
	Thanda mohinah tae, mass murijah aru bimar lagijah khan sai thakibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Litter should be changed periodically and provide clean drinking water and change the water every day. Diseased and dead birds should be immediately removed
	Paikhenah khan somoi tae sapha kurithakibi. Hudai pani bodli kuribi aru sapha pani khelabi. Bimar aru morijah khan joldi para hatailobi



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of DIMAPUR(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	0	23	10	85	46	5.6	68	1
2020-02-02	0	23	10	81	45	6.1	109	0
2020-02-03	0	24	9	56	29	5.4	105	0
2020-02-04	0	22	10	67	34	6.0	112	0
2020-02-05	0	23	10	66	33	6.1	109	1

Weather Summary/Alert:

Probability of no rainfall the coming days Etia ahibleh din tho pani nikiriboleh pareh

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh
--

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Monitor timely for storage pest like moth and weevil infestation
	STORAGE: Dhah ghor tae pukha moth aru weevil ase nah nai somoi tae sai thakibi
GREEN GRAM	STORAGE: Store in gunny bags and kept in clean, ventilated place

Crop(Varieties)	Crop Specific Advisory
	STORAGE : Bijon guti khan bosta tae gusaighenah sapha aru hawa thaka jagah tae rakhibi
RAPESEED	POST HARVEST STAGE: Cleaned seed must be dried in the sun for four to five days
	POST HARVEST STAGE:Sapha bjon guti khan dhup tae 4-5 din sukhabo lageh
MAIZE	HARVESTING STAGE: Dry the cobs immediately before threshing and kernels should be dried sufficiently before storage.
	HARVESTING STAGE: Guti nulua agote makoi bhal parah sukhabo aru guti ulaighenah ghor tae narakha agotae bi bhal para sukhabo
SESAME (GINGELLY/TIL)	STORAGE: Winnow the seeds and dry in the sun for 3 days. Stir once in 3 hours to give uniform drying. Collect the seeds and store in gunnies.
	STORAGE: Guti khan sapha kuri aru dhup tae 3 din sukhabo. Guti sop bhal parah sukhiboleh tin ghonda tae ekbar helabi. Guti the bosta tae rakhibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
LEMON	Using white paint on trunks of tree as it reduces trunk damage
	Ghas biyah nahobo karoneh ghas tae chuna cholabi
FRENCH BEAN	POST HARVEST: French bean are highly perishable, so rapid cooling should be done after harvesting to maintain quality
	POST HARVEST: Quality thakiboleh, harvest kuralogotae thanda jagah tae rakhibi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW	Avoid keeping animals in a damp area, as well as protect them from smoke from fires which are lit to provide warmth. The dampness and smoke increases their chances of contracting pneumonia
	Janwar khan bijah jagah tae narikhibi aru gorom paboleh jui dhunga (smoke) khan bi lagibo nidibi. Etu khan parah pneumonia lagibo pareh
PIG	Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment
	Bimar aru jokom janwar khan joldi parah sailobi, gorom aru bhal jagah tae rakhibi. Veterinarian logotae alusnah kurighenah hi dawai khan khelabi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	If you have drained all the water from the pond, you can clean the bottom of the pond. But do not let your pond becomes too dry. If the bottom becomes too dry, it may crack and will not hold water well when you are ready to fill the pond with water again
	Pukhuri parah pani purah khali kurighenah thakileh, pukhuri nichea sapha kurilobi. Huilobi pukhuri bishi sukha hobo nidibi. Bishi sukha hoileh mathi phatijai aru pani dhali somoi tae pani jamah hui najai

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	It is therefore of major importance to maintain high hygienic and sanitary measures by good management to avoid early exposure of young chicken to diseases. Chutu murgi bacha khan keh bimar nalagibo karoneh sapha aru thik nium parah palibi



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of KIPHIRE(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	1	22	10	83	43	3.3	64	1
2020-02-02	1	24	9	80	24	3.4	56	0
2020-02-03	0	22	8	56	21	4.1	26	0
2020-02-04	0	24	10	68	20	3.0	26	0
2020-02-05	0	25	10	54	17	3.3	56	0

Weather Summary/Alert:

Probability of light rain fall the coming few days
Etia ahiboleh din tho pani olop kiribolhe dikaise

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker
Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh.

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed
Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : Rice should be stored in paddy form rather than milled rice as the husk provides some protection against insects
	STORAGE : Dhan tho sob nukhundighenah, kiman lageh eman hi khundibi. Sob tho khundighenah rakhithakileh puka bi lagijabo

Crop(Varieties)	Crop Specific Advisory
MAIZE	LAND PREPARATION: Deep summer ploughing should be done as it helps in exposing resting stages of pests. To protect seeds from soil borne diseases and insect pest, seed treatment is necessary
	LAND PREPARATION: Mathi phuka khan dhup aparah muriboleh, mathi khunda somoi tae olop dhada khundibi. Bijon guti khan mathi lagah bimar parah bachiboleh seed treatment kuribolageh
MUSTARD	POST HARVESTING : Seed must be dried in sun till the moisture content comes down to 8 percent
	POST HARVESTING : Bijon guti khan moisture 8% tae nalamah thok dhup tae sukhabi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CABBAGE	HARVESTING STAGE: Harvested produce should always be stored in shade before packing.
	Harvest kurighenah packing nukurah agotae, thanda jagah tae rakhibi
BANANA	VEGETATIVE STAGE : Timely remove the dried and diseased leaf
	VEGETATIVE STAGE : Somoi tai sukha aru bimar pata khan ulabi
ONION	HARVESTING STAGE : Bulbs should be harvested soon after the colour of leaves changes to slightly yellow and red pigmentation on bulbs develop.
	HARVESTING STAGE : Pata kitia yellow hoboleh suru hobo, etu somoi tae harvest kuri lobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	At birth, pigs require a temperature around 330C. As the suckling pigs gain weight, they begin to tolerate lower temperatures. Provide artificial lightning.
	Gahori bacha jonom hua somoi tae temperature 330C lageh. Etu karoneh electric light bulb khan cholabi. Bacha khan dangor huija hisap tae, thanda tho loboleh parijai.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minute
	Pani lagah temperature komti huileh, mass khan dana komti khai. Dana bishi nidibi. Dhan bishi dia parah pani letra kureh. 2 minutes tae khai khotom kuribo lagah hisap tae hi dana dibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
HEN	Egg laying is reduced in winter, therefore to increase the production the temperature is to be increased bu installing low wattage bulb.
	Thanda tae anda dia komti hoi. Etu karoneh anda bishi dibo karoneh murghi ghor lagah temperature bishi kuribo lageh



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of KOHIMA(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	2	21	10	99	75	4.9	206	2
2020-02-02	0	24	9	98	54	6.0	202	0
2020-02-03	0	22	8	75	41	5.9	202	0
2020-02-04	0	23	9	94	41	5.5	210	0
2020-02-05	0	24	10	89	37	5.6	210	1

Weather Summary/Alert:

Probability of light rainfall tomorrow. Sky will be mainly clear Etia ahiboleh din tho pani olop kiriboleh dikaiase
--

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh.

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: A good storage system should include prevention of moisture reentering the grain after drying
	STORAGE: Dhan ghor tho dhan sukaighenah rakha khan tae moisture nathakibo hisab tae hobo lageh
RAPSEED	HARVESTING STAGE: It should be harvested when the pods turn yellowish.

Crop(Varieties)	Crop Specific Advisory
FIELD PEA	HARVESTING STAGE: Yellow huija somoi tae harvest kurilobi
	STORAGE : Store the seed in dry bins or in bags kept on wooden racks under cool and dry conditions.
	STORAGE : Bijon guti khan sukha daba aru bags tae jusaighenah khuri lagah racks khan tae rakhibi aru rakha jagah tho dhanda sukha hobo lageh

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	Keep the soil loose by harrowing from time to time
	Somoi somoi tae kol ghas kinar mathi khan dhilah kurighenah rakhibi
CABBAGE	HARVESTING STAGE: Cabbage should be harvested promptly when the heads are firm and mature
	HARVESTING STAGE: Kobi lagah matha dangor huighenah taan hua somoi tae harvest kurilobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	CSF (Classical Swine Fever) may occur in pig at all stages. It is a viral flu and highly contagious. Clinical signs include high fever, skin lesions, convulsions, constipation followed by diarrhoea and vomiting, less appetite. Once the symptoms is noticed, immediately informed the nearby veterinary doctor.
	Chutu dangor gahori sob keh CSF (Classical Swine Fever) lagibo pareh. Etu tho viral flu aru ekta parah ekta keh laghijah bimari ase. Bukhar bishi hua, chamra tae khujuli, dana bhal para nakha, peth jah aru vomiting khan hoi. Etu khan dikhileh, joldi parah osor tae thaka veterinary doctor lagah modot lobi.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Discourage entry of birds and other animals into the pond
	Chirya aru janwar khan pukhuri tae gushiboleh nidibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
HEN	Egg laying is reduced in winter, therefore to increase the production the temperature is to be increased
	Thanda tae anda dia komti hoi. Etu karoneh anda bishi dibo karoneh murgi ghor lagah temperature bishi kuribo lageh



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of LONGLENG(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	1	23	10	93	53	4.8	109	2
2020-02-02	0	25	10	85	40	5.0	135	0
2020-02-03	0	24	9	83	37	5.3	158	0
2020-02-04	0	24	10	79	35	5.0	124	0
2020-02-05	0	25	10	71	32	5.2	124	1

Weather Summary/Alert:

Probability of light shower tomorrow. Sky will be mainly clear

Kali olop pani kiribolhe pareh

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker

Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh.

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed

Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests
	STORAGE: Pukha khan parah bachibo karoneh, dhan rakha somoi tae neem pata logotae rakhibi

Crop(Varieties)	Crop Specific Advisory
RAPESEED	THRESHING: Cleaned seed must be dried in the sun for four to five days or till the moisture content comes down to 8 percent.
	THRESHING: Bijon guti khan moisture 8% tae nalamah thok dhup tae sukhabi
TOMATO	Clean and sanitize harvest containers or bins at least weekly, more often if needed, to remove sand, grit, dirt and other residue
	Clean and sanitize harvest containers or bins at least weekly, more often if needed, to remove sand, grit, dirt and other residue

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	HARVESTING STAGE: 2 TO 3 pickings should be done at 4-5 days interval
	HARVESTING STAGE: 2 bar parah 3 bar nishinah shingilobi aru shinga majotae 4-5 din charighenah hi shingibi
MANDARIN ORANGE	Harvested fruits should be kept under proper storage condition to ensure that there is no change in the quality even after months of storage
	Harvest kurighenah olop mohinah thok biya habobo karoneh bhal parah rakhibi
CABBAGE	HARVESTING STAGE: Cabbage should be harvested promptly when the heads are firm and mature
	HARVESTING STAGE: Kobi lagah matha dangor huighenah taan hua somoi tae harvest kurilobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Vaccinate all newly born piglets against swine fever at the age of 3 months followed by annual booster dose
	Gahori bacha khan 3 months tae swine fever karoneh vaccinate kurilobi aru etu pichetae bi mari thakibi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Check for any fish disease break out or fish mortality during winter months
	Thanda mohinah tae, mass murijah aru bimar lagijah khan sai thakibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Litter should be changed periodically and provide clean drinking water and change the water every day. Diseased and dead birds should be immediately removed
	Paikhenah khan somoi tae sapha kurithakibi. Hudai pani bodli kuribi aru sapha pani khelabi. Bimar aru morijah khan joldi para hatailobi



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of MOKOKCHUNG(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	1	23	10	96	61	4.0	135	2
2020-02-02	0	24	10	94	49	4.5	154	0
2020-02-03	0	23	9	82	40	4.9	135	0
2020-02-04	0	23	10	90	42	4.6	161	0
2020-02-05	0	25	10	82	39	4.6	154	1

Weather Summary/Alert:

Probability of light rainfall tomorrow. Sky will be mainly clear

Etia ahiboleh din tho pani olop kiribolhe dikaiase

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker

Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed

Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests
	STORAGE: Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests

Crop(Varieties)	Crop Specific Advisory
RAPESEED	THRESHING: Cleaned seed must be dried in the sun for four to five days or till the moisture content comes down to 8 percent
	THRESHING: Bijon guti khan moisture 8% tae nalamah thok dhup tae sukhabhi
TOMATO	Clean and sanitize harvest containers or bins at least weekly, more often if needed, to remove sand, grit, dirt and other residue
	Tomato harvest kura daba khan letra bhalu nathakiboleh, somoi tae sapha kuri thakibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	Tomato harvest kura daba khan letra bhalu nathakiboleh, somoi tae sapha kuri thakibi
	HARVESTING STAGE: 2 bar parah 3 bar nishinah shingilobi aru shinga majotae 4-5 din charighenah hi shingibi
ORANGE	Harvested fruits should be kept under proper storage condition to ensure that there is no change in the quality even after months of storage
	Harvest kurighenah olop mohinah thok biya habobo karoneh bhal parah rakhibi
CABBAGE	HARVESTING STAGE: Cabbage should be harvested promptly when the heads are firm and mature
	HARVESTING STAGE: Kobi lagah matha dangor huighenah taan hua somoi tae harvest kurilobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Vaccinate all newly born piglets against swine fever at the age of 3 months followed by annual booster dose
	Gahor bacha khan 3 months tae swine fever karoneh vaccinate kurilobi aru etu pichetae bi mari thakibi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Gahor bacha khan 3 months tae swine fever karoneh vaccinate kurilobi aru etu pichetae bi mari thakibi
	Thanda mohinah tae, mass murijah aru bimar lagijah khan sai thakibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Litter should be changed periodically and provide clean drinking water and change the water every day. Diseased and dead birds should be immediately removed
	Paikhenah khan somoi tae sapha kurithakibi. Hudai pani bodli kuribi aru sapha pani khelabi. Bimar aru morijah khan joldi para hatailobi



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of MON(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	0	24	10	88	56	4.8	154	2
2020-02-02	0	24	9	82	46	5.1	154	0
2020-02-03	0	24	9	78	42	5.3	154	0
2020-02-04	0	24	9	75	38	5.4	158	0
2020-02-05	0	25	10	69	36	5.6	154	1

Weather Summary/Alert:

Probability of no rainfall the coming days, the sky will be mainly clear
Etia ahiboleh din tho pani nikiriboh leh pareh, hoilebi din tho sabha nohobo

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker
Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed
Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Cleanliness should be maintained in the storage area
	Dhan rakha jakha tho sabha rakhibi
RAPSEED	HARVESTING STAGE: The crop is liable to shattering, hence should be harvested just before the pods open in order to avoid losses.

Crop(Varieties)	Crop Specific Advisory
	HARVESTING STAGE: Bhangighenah guti khan gigijabo, etu nuhua agotae harvest kurilobi
FIELD PEA	Green pods should be harvested while just short of reaching maturity. Multiple picking like, 4 to 5 picking should be done within the 2 to 10 days. 2 to 10 days dhe apni khan charr par pass bar tho koti dho jikipo lhe pareh
GARLIC	STORAGE: Garlic with dried leaves can be stored by hanging in well-ventilated rooms.
	STORAGE: Losoon pata bhal para sukijah khan hawa aha jah kurah room tae hang kurighenah rakhibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
POTATO	HARVESTING STAGE: The harvested potatoes should be surface dried and kept in heaps for 10-15 days in shade for curing of skin. The tubers should not be exposed to direct sunlight as they become green
	Alu tho harvest khorabiche dhe direct doob lakha nichhe dhe narakhibi
ORANGE	To kill the trunk borer grubs, clean the bored holes of the infested trunk with iron wire and insert a cotton swab soaked in the mixture of kerosene and water(1:1) and plug with mud
	Ghas lakha korithe pukha khan daka tho, wire parah sabha khoribi etu biche de cotton kerosene aru pani mixture (1:1) DHE BICHABI ETU BICHE THE jenta the kusaidhebi. Mati para jenta tho pon koribi
ONION	STORAGE: Onions should not be stored unless adequately dried either in the field or by artificial means
	Piyas tho pal para sukhi khena hai rakhibi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Good housekeeping practices such as removing manure, washing down pens, disinfecting and generally cleaning up the facility are essential to maintaining a healthy herd
	Gauri bimar nahokina gao pal hobo garoneh, ghor dho sabha rakhibo lageh

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Discourage entry of birds and other animals into the pond
	Chirya aru janwar khan pukhuri tae gushiboleh nidibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
HEN	You can improve the amount of meat and eggs by providing better housing for birds and good feed and water
	Murgi karoneh ghor bhal aru bhal dana pani khela parah murgi gao dangor hoi aru anda bishi dae.



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of PEREN(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	0	23	10	89	49	3.6	124	2
2020-02-02	0	24	10	79	40	4.6	158	0
2020-02-03	0	23	8	57	30	4.8	116	0
2020-02-04	0	23	9	76	28	4.7	112	0
2020-02-05	0	24	10	70	27	4.7	112	2

Weather Summary/Alert:

Probability of no rainfall the coming days
Etia ahibleh din tho pani nikiriboleh pareh

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker
Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed
Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Monitor timely for storage pest like moth and weevil infestation
	STORAGE: Dhah ghor tae pukha moth aru weevil ase nah nai somoi tae sai thakibi
GREEN GRAM	STORAGE: Store in gunny bags and kept in clean, ventilated place

Crop(Varieties)	Crop Specific Advisory
	STORAGE : Bijon guti khan bosta tae gusaighenah sapha aru hawa thaka jagah tae rakhibi
RAPESEED	POST HARVEST STAGE: Cleaned seed must be dried in the sun for four to five days
	POST HARVEST STAGE:Sapha bjon guti khan dhup tae 4-5 din sukhabo lageh
MAIZE	HARVESTING STAGE: Dry the cobs immediately before threshing and kernels should be dried sufficiently before storage.
	HARVESTING STAGE: Guti nulua agote makoi bhal parah sukhabi aru guti ulaighenah ghor tae narakha agotae bi bhal para sukhabi
SESAME (GINGELLY/TIL)	STORAGE: Winnow the seeds and dry in the sun for 3 days. Stir once in 3 hours to give uniform drying. Collect the seeds and store in gunnies
	STORAGE: Winnow the seeds and dry in the sun for 3 days. Stir once in 3 hours to give uniform drying. Collect the seeds and store in gunnies

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
LEMON	Using white paint on trunks of tree as it reduces trunk damage
	Ghas biyah nahobo karoneh ghas tae chuna cholabi
FRENCH BEAN	POST HARVEST: French bean are highly perishable, so rapid cooling should be done after harvesting to maintain quality
	POST HARVEST: Quality thakiboleh, harvest kuralogotae thanda jagah tae rakhibi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
COW	Avoid keeping animals in a damp area, as well as protect them from smoke from fires which are lit to provide warmth. The dampness and smoke increases their chances of contracting pneumonia
	Janwar khan bijah jagah tae narikhibi aru gorom paboleh jui dhunga (smoke) khan bi lagibo nidibi. Etu khan parah pneumonia lagibo pareh
PIG	Sick or injured animals should be attended to immediately, preferably move to a warm comfortable environment. Medications should be used only on the advice of a veterinarian, following recommended treatment
	Bimar aru jokom janwar khan joldi parah sailobi, gorom aru bhal jagah tae rakhibi. Veterinarian logotae alusnah kurighenah hi dawai khan khelabi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	If you have drained all the water from the pond, you can clean the bottom of the pond. But do not let your pond becomes too dry. If the bottom becomes too dry, it may crack and will not hold water well when you are ready to fill the pond with water again
	Pukhuri parah pani purah khali kurighenah thakileh, pukhuri nichea sapha kurilobi. Huilobi pukhuri bishi sukha hobo nidibi. Bishi sukha hoileh mathi phatijai aru pani dhalia somoi tae pani jamah hui najai

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
CHICK	It is therefore of major importance to maintain high hygienic and sanitary measures by good management to avoid early exposure of young chicken to diseases. Chutu murgi bacha khan keh bimar nalagibo karoneh sapha aru thik nium parah palibi



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of PHEK(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	3	20	10	89	58	3.2	206	2
2020-02-02	0	23	9	84	33	3.7	216	0
2020-02-03	0	21	8	71	30	3.6	292	0
2020-02-04	0	24	9	76	27	3.0	252	0
2020-02-05	0	24	10	68	23	2.8	246	1

Weather Summary/Alert:

Probability of light rainfall tomorrow. Sky will be mainly clear kali olop pani kiribolhe pareh
--

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh
--

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: A good storage system should include prevention of moisture reentering the grain after drying
	STORAGE: Dhan ghor tho dhan sukaighenah rakha khan tae moisture nathakibo hisab tae hobo lageh
RAPESEED	HARVESTING STAGE: It should be harvested when the pods turn yellowish.

Crop(Varieties)	Crop Specific Advisory
FIELD PEA	HARVESTING STAGE: Yellow huija somoi tae harvest kurilobi
	STORAGE : Store the seed in dry bins or in bags kept on wooden racks under cool and dry conditions
	STORAGE : Bijon guti khan sukha daba aru bags tae jusaighenah khuri lagah racks khan tae rakhibi aru rakha jagah tho dhanda sukha hobo lageh

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
BANANA	STORAGE : Bijon guti khan sukha daba aru bags tae jusaighenah khuri lagah racks khan tae rakhibi aru rakha jagah tho dhanda sukha hobo lageh
	Somoi somoi tae kol ghas kinar mathi khan dhilah kurighenah rakhibi
CABBAGE	HARVESTING STAGE: Cabbage should be harvested promptly when the heads are firm and mature
	HARVESTING STAGE: Kobi lagah matha dangor huighenah taan hua somoi tae harvest kurilobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	CSF (Classical Swine Fever) may occur in pig at all stages. It is a viral flu and highly contagious. Clinical signs include high fever, skin lesions, convulsions, constipation followed by diarrhoea and vomiting, less appetite. Once the symptoms is noticed, immediately informed the nearby veterinary doctor.
	Chutu dangor gahori sob keh CSF (Classical Swine Fever) lagibo pareh. Etu tho viral flu aru ekta parah ekta keh laghijah bimari ase. Bukhar bishi hua, chamra tae khujuli, dana bhal para nakha, peth jah aru vomiting khan hoi. Etu khan dikhileh, joldi parah osor tae thaka veterinary doctor lagah modot lobi.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Discourage entry of birds and other animals into the pond
	Chiryra aru janwar khan pukhuri tae gushiboleh nidibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
HEN	Egg laying is reduced in winter, therefore to increase the production the temperature is to be increased
	Thanda tae anda dia komti hoi. Etu karoneh anda bishi dibo karoneh murgi ghor lagah temperature bishi kuribo lageh



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of TUENSANG(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	2	23	10	97	51	4.1	112	2
2020-02-02	1	25	10	90	32	4.0	135	0
2020-02-03	0	23	9	85	32	4.3	112	0
2020-02-04	0	24	10	83	31	3.9	112	0
2020-02-05	0	26	11	70	25	4.0	109	0

Weather Summary/Alert:

Probability of light shower the coming days few days

Etia ahiboleh din tho pani olop kiribolhe dikaise

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker

Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh.

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed

Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE : Rice should be stored in paddy form rather than milled rice as the husk provides some protection against insects
	STORAGE : Dhan tho sob nukhundighenah, kiman lageh eman hi khundibi. Sob tho khundighenah rakhithakileh puka bi lagijabo

Crop(Varieties)	Crop Specific Advisory
MAIZE	LAND PREPARATION: Deep summer ploughing should be done as it helps in exposing resting stages of pests. To protect seeds from soil borne diseases and insect pest, seed treatment is necessary
	LAND PREPARATION: Mathi phuka khan dhup aparah muriboleh, mathi khunda somoi tae olop dhada khundibi. Bijon guti khan mathi lagah bimar parah bachiboleh seed treatment kuribolageh
MUSTARD	POST HARVESTING : Seed must be dried in sun till the moisture content comes down to 8 percent
	POST HARVESTING : Bijon guti khan moisture 8% tae nalamah thok dhup tae sukhabi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
CABBAGE	HARVESTING STAGE: Harvested produce should always be stored in shade before packing.
	Harvest kurighenah packing nukurah agotae, thanda jagah tae rakhibi
BANANA	VEGETATIVE STAGE : Timely remove the dried and diseased leaf
	VEGETATIVE STAGE : Somoi tai sukha aru bimar pata khan ulabi
ONION	HARVESTING STAGE : Bulbs should be harvested soon after the colour of leaves changes to slightly yellow and red pigmentation on bulbs develop.
	HARVESTING STAGE : Pata kitia yellow hoboleh suru hobo, etu somoi tae harvest kuri lobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	At birth, pigs require a temperature around 33 degree C. As the suckling pigs gain weight, they begin to tolerate lower temperatures. Provide artificial lightning
	Gahori bacha jonom hua somoi tae temperature 330C lageh. Etu karoneh electric light bulb khan cholabi. Bacha khan dangor huija hisap tae, thanda tho loboleh parijai.

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	As the water temperature of your pond falls, the fish will require less food. Never over feed your fish. Excess food will pollute the pond water, so only feed amounts that will be completely consumed by your fish within 2 minutes.
	Pani lagah temperature komti huileh, mass khan dana komti khai. Dana bishi nidibi. Dhan bishi dia parah pani letra kureh. 2 minutes tae khai khotom kuribo lagah hisap tae hi dana dibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICKEN	Egg laying is reduced in winter, therefore to increase the production the temperature is to be increased bu installing low wattage bulb.
	Thanda tae anda dia komti hoi. Etu karoneh anda bishi dibo karoneh murghi ghor lagah temperature bishi kuribo lageh



Gramin Krishi Mausam Sewa
India Meteorological Department
ICAR Complex Branch
Umiam, Meghalaya



Agromet Advisory Bulletin

Date : 31-01-2020

Weather Forecast of WOKHO(Nagaland) Issued On : 2020-01-31(Valid Till 08:30 IST of the next 5 days)

Date (y-m-d)	Rainfall (mm)	Tmax (°C)	Tmin (°C)	RH I (%)	RH II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2020-02-01	0	24	10	98	55	4.3	109	2
2020-02-02	0	23	10	98	53	4.7	109	0
2020-02-03	0	24	9	73	36	4.8	112	0
2020-02-04	0	23	9	92	43	5.0	112	0
2020-02-05	0	24	9	86	39	4.9	110	1

Weather Summary/Alert:

Probability of no rainfall the coming days. Sky will be mainly clear

Etia ahibleh din tho pani nikiribolhe pareh

General Advisory:

Give the animals (cattle) 30 – 50 grams of mineral mixture along with their feed on a daily basis. This increases milk productivity and enhances the animal's immunity. Animals must be dewormed, using the correct dose of medicines, after due consultation from the veterinarian / animal health worker

Janwar keh hudai danah dia tae 30-50 grams mineral mixture milaighenah dibi. Etu parah janwar khan bimar parah bishibhak bachighenah rakhe aru doot dai bi bishi hoi. Veterinarian nahoileh animal health worker khan logotae alusnah kurighenah janwar khan keh kechu dawai khelabo lageh

SMS Advisory:

In case of diarrhea in poultry, Always ensure that your birds have fresh clean drinking water, fresh good quality feed

Murkhi khan peth bishi jaa somai tae, hudai pani diya tho sapha hoi nah nahoi sabo lageh aru dana bi thik jaga teh rakha laga aru biya nuha khelabo lageh

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	STORAGE: Paddy grains can be mixed with neem leaves at the time of storage to reduce storage pests
	STORAGE: Pukha khan parah bachibo karoneh, dhan rakha somoi tae neem pata logotae rakhibi

Crop(Varieties)	Crop Specific Advisory
RAPESEED	THRESHING: Cleaned seed must be dried in the sun for four to five days or till the moisture content comes down to 8 percent
	THRESHING: Bijon guti khan moisture 8% tae nalamah thok dhup tae sukhabhi
TOMATO	Clean and sanitize harvest containers or bins at least weekly, more often if needed, to remove sand, grit, dirt and other residue
	Tomato harvest kura daba khan letra bhalu nathakiboleh, somoi tae sapha kuri thakibi

Horticulture Specific Advisory:

Horticulture (Varieties)	Horticulture Specific Advisory
FRENCH BEAN	HARVESTING STAGE: 2 TO 3 pickings should be done at 4-5 days interval
	HARVESTING STAGE: 2 bar parah 3 bar nishinah shingilobi aru shinga majotae 4-5 din charighenah hi shingibi
MANDARIN ORANGE	Harvested fruits should be kept under proper storage condition to ensure that there is no change in the quality even after months of storage
	Harvest kurighenah olop mohinah thok biya habobo karoneh bhal parah rakhibi
CABBAGE	HARVESTING STAGE: Cabbage should be harvested promptly when the heads are firm and mature
	HARVESTING STAGE: Kobi lagah matha dangor huighenah taan hua somoi tae harvest kurilobi

Livestock Specific Advisory:

Livestock(Varieties)	Livestock Specific Advisory
PIG	Vaccinate all newly born piglets against swine fever at the age of 3 months followed by annual booster dose
	Gahor bacha khan 3 months tae swine fever karoneh vaccinate kurilobi aru etu pichetae bi mari thakibi

Fisheries Specific Advisory:

Fisheries(Varieties)	Fisheries Specific Advisory
COMMON FISH	Check for any fish disease break out or fish mortality during winter months.
	Thanda mohinah tae, mass murijah aru bimar lagijah khan sai thakibi

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
FOWL	Litter should be changed periodically and provide clean drinking water and change the water every day. Diseased and dead birds should be immediately removed
	Paikhenah khan somoi tae sapha kurithakibi. Hudai pani bodli kuribi aru sapha pani khelabi. Bimar aru morijah khan joldi para hatailobi