


STATE COMPOSITE AGRO-MET ADVISORY BULLETIN FOR THE STATE OF SIKKIM



PERIOD: 11th MAY to 15th MAY'2019

IMD, GANGTOK IN COLLABORATION WITH ICAR, GANGTOK, FSAD AND HCCD, GANGTOK





ISSUED ON TUESDAY 10th MAY'2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for NORTH-DISTRICT for next five days

Parameter/Date	11/05/2019	12/05/2019	13/05/2019	14/05/2019	15/05/2019
Rainfall (mm)	Moderate rainfall	Light rainfall	Light Rainfall	Light Rainfall	Moderate rainfall
Max Temp (⁰ C)	28	28	28	28	27
Min Temp (⁰ C)	18	18	18	18	18
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	95	92	94	91	95
Min RH (%)	85	75	78	78	87
Wind Speed (KmpH)	3	3	3	2	2
Wind Direction (deg)	E	S	S	W	SE

Northerly- N, North-Easterly- E, South-Easterly- S-E, South-Westerly- S, South-Westerly- N, Westerly- W, North-westerly- N-W.

Weather for preceding week					
Date	Rainfall	Max. Temp	Min. Temp		
	(mm)	(°C)	(°C)		
07/05/2019	000	28.0	08.2		
08/05/2019	000	27.0	09.1		
09/06/2019	000	28.0	08.2		

- > Probability of light rainfall
- Max temp may be varying from 27-28°C, whereas, Min temp may vary from 18°C
- > Sky will remain generally cloudy
- Max RH is likely to be 91-95 %, whereas, Min RH may vary from 75-85 %
- > Wind speed may likely vary from 2-3 km/hr.
- > Wind direction may blow from easterly to southerly and then westerly to finally southeasterly
- > Weekly cumulative rainfall=49 mm
- > Heavy Rain likely to occur at one or two places over the district on Tuesday.

Value Added forecast for EAST-DISTRICT for next five days

Parameter/Date	11/05/2019	12/05/2019	13/05/2019	14/05/2019	15/05/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	22	21	21	21	21
Min Temp (⁰ C)	17	17	17	16	16
Sky condition	Cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	95	94	92	94	95
Min RH (%)	85	84	80	85	87
Wind Speed (KmpH)	3	4	4	2	3
Wind Direction (deg)	NE	NE	NE	NE	NE

Northerly- N, North-Easterly- E, South-Easterly- S-E, South-Westerly- S, South-Westerly- S, Westerly- W, North-westerly- N-W.

	Weather for preceding week				
Date	Rainfall	Max. Temp	Min. Temp		
	(mm)	(°C)	(°C)		
07/05/2019	003	22.8	15.0		
08/05/2019	023	20.3	15.7		
09/06/2019	025	21.6	16.7		

- > Probability of light rainfall
- Max temp may be varying from 21-22°C, whereas, Min temp may vary from 16-17 °C
- > Sky will remain generally cloudy
- Max RH is likely to be 92-95 %, whereas, Min RH may vary from 80-85 %
- > Wind speed may likely vary from 2-4 km/hr.
- > Wind direction may blow from northeasterly
- > Weekly cumulative rainfall= 38 mm

Value Added forecast for SOUTH-DISTRICT for next five days

Parameter/Date	11/05/2019	12/05/2019	13/05/2019	14/05/2019	15/05/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	28	28	28	27	27
Min Temp (⁰ C)	18	17	17	17	17
Sky condition	Company lley alouder	Generally	Comovaller aloreder	C II1 I	Generally cloudy
	Generally cloudy	cloudy	Generally cloudy	Generally cloudy	
Max RH (%)	91	90	90	91	92
Min RH (%)	70	68	65	70	76
Wind Speed (KmpH)	3	4	3	2	3
Wind Direction (deg)	NE	NE	NE	NE	NE

Northerly- N, North-Easterly- E, South-Easterly- S, South-Westerly- S, South-Westerly- N, Westerly- N, Wester

	Weather for preceding week				
Date Rainfall Max. Temp Min. Temp					
	(mm)	(°C)	(°C)		
07/05/2019	005	27.5	15.5		
08/05/2019	000	N/A	15.0		
09/06/2019	000	28.0	N/A		

- There will be chances of light rainfall
- > Max temp may be varying from 27-28°C, whereas, Min temp may vary from 17-18°C
- > Sky will remain partially clear
- > Max RH is likely to be 85-90 %, whereas, Min RH may var from 64-70 %
- > Wind speed may likely vary from 3-4 km/hr.
- > Wind direction may blow from north easterly
- > Weekly cumulative rainfall= 24 mm

Value Added forecast for WEST-DISTRICT for next five days

Parameter/Date	11/05/2019	12/05/2019	13/05/2019	14/05/2019	15/05/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	25	24	25	25	24
Min Temp (⁰ C)	17	17	17	17	16
Sky condition	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy	Generally cloudy
Max RH (%)	92	90	90	92	92
Min RH (%)	80	75	75	80	81
Wind Speed (KmpH)	3	4	3	2	3
Wind Direction (deg)	NW	NW	NW	S	NW

Northerly- N, North-Easterly- E, South-Easterly- S, South-Westerly- S, South-Westerly- N, Westerly- N, North-westerly- N, North

	Weather for preceding week				
Date Rainfall Max. Temp Min. Temp					
	(mm)	(°C)	(°C)		
07/05/2019	N/A	24.5	14.7		
08/05/2019	003	22.6	16.5		
09/06/2019	005	24.6	17.0		

- > Probability of light rainfall
- > Max temp may be varying from 24-25°C, whereas, Min temp may vary from 16-17°C
- > Sky will remain generally cloudy
- Max RH is likely to be 90-92 %, whereas, Min RH may var from 75-81 %
- > Wind speed may likely vary from 3-4 km/hr.
- > Wind direction may blow from Northwesterly to southerly and finally northwesterly
- > Weekly cumulative rainfall= 31 mm

AGRO-METEREOLOGICAL ADVISORIES

General Advisories for all the crops/ livestock

- Bunding at terrace should be done to reduce nutrient losses due to heavy rainfall
- Keep ready the field for paddy cultivation
- Jalkund should be dug out for rain water harvesting.
- **▶** Spraying of plant protection bio pesticides on bright sunny day for insect and disease management is recommended.
- Maintain strict hygiene and sanitation measures in poultry farm.
- **▶** Strategic supplementation of vitamins and mineral mixture should provide to livestock to improve the productive performance.
- Prove green grass to the livestock to feed as much as possible

		,					
Name of the	Stage Cultural practices/			Agro-Met Advisories			
crop/Animal		Pest/ Dis					
	Agricultural and Horticultural crops:						
Maize	Knee heigh	at stage	Army worm	 ♦ Field should be weed free without damaging the main crop while uprooting the weeds. ♦ Earthing up should be done. ♦ Apply bio fertilizers @25 kg/ha and vermicompost @1.5 t/ha. ♦ Foliar spray of neem based formulation @ 0.5% or 5 ml/lit. of water at weekly interval 			
Rice	Field prepa	aration		Start initial ploughing			
Ginger/ Turmeric Large cardamom	Field preparation Sowing Planting			 Apply bio mulch @10 t/ha to cover the ridges and furrows of the field A 2 years old nursery suckers should be planted in the main field A spacing of 1.8×1.8 m² should be maintained. Young white coloured roots should not be destroy while planting, while old roots should trimmed out. 			

Early kharif vegetables	Vegetative growth		Weeding and intercultural operation should be done
Tomato/Red cherry pepper/Brinjal	Vegetative	Blight	 Plants should be keep weed free and avoid water stagnation Spraying of COC @2-3 ml/lit. at weekly interval
Sikkim Mandarin	Planting/fruiting		 Keep the orchard weed free Digging the pit of 3 cubic feet for transplanting new saplings Top soil of the pit should be mixed with FYM @20 kg/pit
	Animal Sciences	and livestock	management
Tibetan Sheep			 Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended.
Yak			 Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended.
Sheep and Goat Cattle			 Trimming of hair is recommended. Farmers are advised to look for heat sign especially during early morning or late evening and ensure breeding of the dairy cattle 12 hr after observation of heat. Breeding of the sheep/goat should be done 24-30 hr after observation of heat. Feeding of concentrate feed along with jaggery, mineral mixture and salt. Periodic application of fly repellents and/or netting of dung pit are recommended. Balanced feeding of dairy animals
			and allow calf suckling immediately after parturition. Colostrum (first three days milk) feeding to new born calf for first three days of life should be ensured. Full hand milking should be practiced. Frequent incomplete milking at interval of 4hrs should be practiced after parturition for minimum 1 week. Afterwards complete milking in a day

			is recommended.
Poultry		Coccidiosis, Ranikhet disease	 16 hrs of light (natural and electrical) should be provided to adult birds for optimum egg production. Frequent raking and liming of deep litter materials. Inclusion of turmeric powder @ 10g/kg feed for prevention of coccidiosis. Maintenance of brooder temperature for chicks with the help of electrical bulbs for initial one month. Ranikhet vaccination at 5-7 days and at 28-30 days of age. Calcium/shell grit and multi vitamin supplementation is strongly recommended to optimize egg production. Spraying of ectoparasiticides viz., deltamethrin/cypermethrin @ 1% v/v in water.
Pig	All stages	Swine fever	 Keep floor dry to avoid skin diseases. Weaning is recommended at the age of 8 weeks for crossbreds and 10 weeks for indigenous pigs. Swine fever vaccine at the age of 3 months is recommended. Critical heat detection in early morning and late evening is recommended. Feed 5-10% green fodder of their diet.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim Designation Name **Department** Dr. R. K. Avasthe **Soil Science Joint Director** Dr. Raghavendra Singh Senior scientist and Head(I/C) Agronomy Dr. P. K. Pathak Animal science SMS Dr. J.K. Singh Plant Breeding SMS Dr. Manoj Kumar Agronomy SMS Sh. Boniface Lepcha SMS Horticulture SMS Agril. Meteorology Sh. Santanu Das Metrological observer Biochemistry Miss Tirtha Kumari Chettri