


STATE COMPOSITE AGRO-MET ADVISORY BULLETIN FOR THE STATE OF SIKKIM PERIOD: 18th MAY to 22nd MAY'2019



IMD, GANGTOK IN COLLABORATION WITH *ICAR, GANGTOK, FSAD AND HCCD, GANGTOK*





ISSUED ON TUESDAY 17th MAY'2019

Meteorological Sub-division of Sikkim

As per India Meteorological Department's classification, Sikkim is a part of Met Sub-Division of Sub Himalayan West Bengal & Sikkim (SHWB & SKM) and is divided into four (4) Districts:

- 1) North District with HQ at Mangan
- 2) East District with HQ at Gangtok
- 3) West District with HQ at Gyalsing
- 4) South District with HQ at Namchi

Issued by: Santanu Das, SMS (Agril. Meteorology), ICAR-KVK, East Sikkim, Ranipool

Value Added forecast for NORTH-DISTRICT for next five days

Parameter/Date	18/05/2019	19/05/2019	20/05/2019	21/05/2019	22/05/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	28	29	29	29	28
Min Temp (⁰ C)	18	18	18	18	17
Sky condition	Generally cloudy	Generally cloudy	Partially clear	Partially clear	Generally cloudy
Max RH (%)	92	92	88	90	90
Min RH (%)	75	75	72	74	74
Wind Speed (KmpH)	003	004	003	003	002
Wind Direction (deg)	SW	S	SW	SW	SW

Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E, Southerly- S, South-Westerly- S-W, Westerly-W, North-westerly- N-W.

Weather for preceding week			
Date	Rainfall	Max. Temp	Min. Temp
	(mm)	(°C)	(°C)
14/05/2019	011	26.0	08.1
15/05/2019	003	20.0	08.0
16/06/2019	007	23.0	09.0

- >> Probability of light rainfall
- > Max temp may be varying from 28-29°C, whereas, Min temp may vary from 17-18°C
- >> Sky will remain generally cloudy
- >> Max RH is likely to be 88-92 %, whereas, Min RH may vary from 72-75 %
- >> Wind speed may likely vary from 2-4 km/hr.
- > Wind direction may blow from southeasterly to southerly , then southerly to southwesterly and finally
- > Weekly cumulative rainfall=42 mm

Value Added forecast for EAST-DISTRICT for next five days

Parameter/Date	18/05/2019	19/05/2019	20/05/2019	21/05/2019	22/05/2019
Rainfall (mm)	Light rainfall	Light rainfall	Light rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	22	23	23	22	22
Min Temp (⁰ C)	16	17	17	16	16
Sky condition	Generally cloudy	Generally cloudy	Partially clear	Partially clear	Generally cloudy
Max RH (%)	92	91	89	89	91
Min RH (%)	76	76	74	75	76
Wind Speed (KmpH)	002	004	003	002	002
Wind Direction (deg)	NE	NE	NE	NE	NE

Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E, Southerly- S, South-Westerly- S-W, Westerly-W, North-westerly- N-W.

	Weather for preceding week					
Date	Rainfall	Max. Temp	Min. Temp			
	(mm)	(°C)	(°C)			
14/05/2019	001	20.7	15.2			
15/05/2019	000	19.5	14.7			
16/06/2019	004	18.8	14.7			

- > Probability of light rainfall
- >> Max temp may be varying from 22-23°C, whereas, Min temp may vary from 16-17 °C
- >> Sky will remain partially clear
- Max RH is likely to be 89-92 %, whereas, Min RH may vary from 74-76 %
- >> Wind speed may likely vary from 2-4 km/hr.
- > Wind direction may blow from northeasterly
- > Weekly cumulative rainfall= 40 mm

Value Added forecast for SOUTH-DISTRICT for next five days

Parameter/Date	18/05/2019	19/05/2019	20/05/2019	21/05/2019	22/05/2019
Rainfall (mm)	No rain	No rain	Light rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	27	28	28	28	27
Min Temp (⁰ C)	17	18	18	18	17
Sky condition	Generally cloudy	Partially clear	Generally cloudy	Partially clear	Partially clear
Max RH (%)	88	88	90	88	88
Min RH (%)	70	70	75	70	70
Wind Speed (KmpH)	003	004	003	003	002
Wind Direction (deg)	NE	NE	NE	NE	NE

Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E, Southerly- S, South-Westerly- S-W, Westerly-W, North-westerly- N-W.

	Weather for preceding week					
Date	Rainfall	Max. Temp	Min. Temp			
	(mm)	(°C)	(°C)			
14/05/2019	000	27.5	12.5			
15/05/2019	000	28.0	13.5			
16/06/2019	011	N/A	15.0			

- > There will be chances of light rainfall
- >> Max temp may be varying from 27-28°C, whereas, Min temp may vary from 17-18°C
- > Sky will remain generally cloudy
- >> Max RH is likely to be 88-90 %, whereas, Min RH may var from 70-75 %
- >> Wind speed may likely vary from 2-4 km/hr.
- > Wind direction may blow from north easterly
- > Weekly cumulative rainfall= 18 mm

Value Added forecast for WEST-DISTRICT for next five days

Parameter/Date	15/05/2019	16/05/2019	17/05/2019	18/05/2019	19/05/2019
Rainfall (mm)	Light rainfall	Light rainfall	No rainfall	Light rainfall	Light rainfall
Max Temp (⁰ C)	24	25	25	25	25
Min Temp (⁰ C)	17	18	18	17	17
Sky condition	Generally cloudy	Generally cloudy	Partially clear	Generally cloudy	Generally cloudy
Max RH (%)	90	90	88	92	91
Min RH (%)	75	75	72	76	75
Wind Speed (KmpH)	004	005	003	003	002
Wind Direction (deg)	NW	NW	NW	NW	NW

Northerly- N, North-Easterly- N-E, Easterly- E, South-Easterly- S-E, Southerly- S, South-Westerly- S-W, Westerly-W, North-westerly- N-W.

	Weather for preceding week					
Date	Rainfall	Max. Temp	Min. Temp			
	(mm)	(°C)	(°C)			
14/05/2019	N/A	22.1	15.0			
15/05/2019	000	21.7	13.9			
16/06/2019	003	21.0	14.2			

- >> Probability of light rainfall
- > Max temp may be varying from 24-25°C, whereas, Min temp may vary from 17-18°C
- > Sky will remain generally cloudy
- Max RH is likely to be 88-92 %, whereas, Min RH may var from 72-76 %
- Wind speed may likely vary from 2-5 km/hr.
- > Wind direction may blow from northwesterly
- > Weekly cumulative rainfall= 26 mm
- > Hailstorm may prevail on Wednesday

AGRO-METEREOLOGICAL ADVISORIES

General Advisories for all the crops/ livestock

- Bunding at terrace should be done to reduce nutrient losses due to heavy rainfall
- Keep ready the field for paddy cultivation
- *Jalkund* should be dug out for rain water harvesting.
- Spraying of plant protection bio pesticides on bright sunny day for insect and disease management is recommended.
- Maintain strict hygiene and sanitation measures in poultry farm.
- Strategic supplementation of vitamins and mineral mixture should provide to livestock to improve the productive performance.
- Prove green grass to the livestock to feed as much as possible

Name of the crop/Animal	Stage	Cultural p Pest/Dis		Agro-Met Advisories
	Agricu	ultural and H	lorticultu	iral crops:
Maize	Knee heigh	t stage	Army worm	 Field should be weed free without damaging the main crop while uprooting the weeds. Earthing up should be done. Apply bio fertilizers @25 kg/ha and vermicompost @1.5 t/ha. Foliar spray of neem based formulation @ 0.5% or 5 ml/lit. of water at weekly interval
Rice	Field prepa	aration		Start initial ploughing
Ginger/ Turmeric	Sowin	ıg		Apply bio mulch @10 t/ha to cover the ridges and furrows of the field
Large cardamom	Planting Value of the second s			 A 2 years old nursery suckers should be planted in the main field A spacing of 1.8×1.8 m² should be maintained. Young white coloured roots should not be destroy while planting, while old roots should trimmed out.

		I			
Early kharif vegetables	Vegetative growth			۲	Weeding and intercultural operation should be done
Tomato/Red cherry pepper/Brinjal	Vegetative		Blight	۲ ۲	Plants should be keep weed free and avoid water stagnation Spraying of COC @2-3 ml/lit. at weekly interval
Sikkim Mandarin	Planting/fruiting			۲	Keep the orchard weed free Digging the pit of 3 cubic feet for transplanting new saplings Top soil of the pit should be mixed with FYM @20 kg/pit
	Animal Sciences	s and li	ivestock	ma	nagement
Tibetan Sheep					Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended.
Yak				 <td>Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended. Trimming of hair is recommended.</td>	Feeding of concentrate feed along with mineral mixture and salt. Hay feeding is recommended. Trimming of hair is recommended.
Sheep and Goat				0	Farmers are advised to look for heat
				٩	sign especially during early morning or late evening and ensure breeding of the dairy cattle 12 hr after observation of heat. Breeding of the sheep/goat should be done 24-30 hr after observation of heat.
				۲	Feeding of concentrate feed along with jaggery, mineral mixture and salt.
				۲	Periodic application of fly repellents and/or netting of dung pit are recommended.
Cattle				•	Balanced feeding of dairy animals and allow calf suckling immediately after parturition. Colostrum (first three days milk) feeding to new born calf for first three days of life should be ensured. Full hand milking should be
				۲	practiced. Frequent incomplete milking at interval of 4hrs should be practiced after parturition for minimum 1 week. Afterwards complete milking in a day

			is recommended.
Poultry		Coccidiosis, Ranikhet disease	 16 hrs of light (natural and electrical) should be provided to adult birds for optimum egg production. Frequent raking and liming of deep litter materials. Inclusion of turmeric powder @ 10g/kg feed for prevention of coccidiosis. Maintenance of brooder temperature for chicks with the help of electrical bulbs for initial one month. Ranikhet vaccination at 5-7 days and at 28-30 days of age. Calcium/shell grit and multi vitamin supplementation is strongly recommended to optimize egg production. Spraying of ectoparasiticides <i>viz.</i>, deltamethrin/cypermethrin @ 1% v/v in water.
Pig	All stages	Swine fever	 Keep floor dry to avoid skin diseases. Weaning is recommended at the age of 8 weeks for crossbreds and 10 weeks for indigenous pigs. Swine fever vaccine at the age of 3 months is recommended. Critical heat detection in early morning and late evening is recommended. Feed 5-10% green fodder of their diet.

Agro-Met Advisory member of ICAR-NOFRI and KVK- East Sikkim					
Name Dr. R. K. Avasthe	Designation Joint Director	Department Soil Science			
Dr. Raghavendra Singh	Senior scientist and Head(I/C)	Agronomy			
Dr. P. K. Pathak	SMS	Animal science			
Dr. J.K. Singh	SMS	Plant Breeding			
Dr. Manoj Kumar	SMS	Agronomy			
Sh. Boniface Lepcha	SMS	Horticulture			
Sh. Santanu Das	SMS	Agril. Meteorology			
Miss Tirtha Kumari Chettri	Metrological observer	Biochemistry			