

## Medicinal Uses of Roselle Plant (*Hibiscus sabdariffa* L.): A Mini Review

K. PURO\*, R. SUNJUKTA, S. SAMIR, S. GHATAK, I. SHAKUNTALA, A. SEN

Received 8.6. 2014, Revised 28.6.2014, Accepted 30.6.2014

### ABSTRACT

The use of herbal extracts and nutritional supplements either as alternative or complementary medicine for treatment of diseases is well documented in various cultures such as Ayurveda in India and traditional Chinese medicine system. Medicinal plants as natural antimicrobial agents are gaining popularity. Roselle plant (*Hibiscus sabdariffa*) Linne (Malvaceae) has been used in folk medicine as a diuretic, mild laxative, and treatment for cardiac and nerve diseases. Herein we discussed some of the recent studies on its various activities.

**Keywords:** Antioxidant, Antihypertensive, Antimicrobial, Anticancerous

### INTRODUCTION

Roselle plant, with an attractive flower believed to be native to Africa, is cultivated in Sudan, India, Malaysia and Taiwan. It is an annual or perennial herb or woody-based sub-shrub, growing to 2-2.5 m tall. The leaves are deeply 3-5 lobed, 8-15 cm long, arranged alternately on the stems. The flowers are 8-10 cm in diameter, white to pale yellow with a dark red spot at the base of each petal, and have a stout fleshy calyx at the base, 1-2 cm wide, enlarging to 3-3.5 cm, fleshy and bright red as the fruit matures. It takes about six months to mature. In some places, the plant is primarily cultivated for the production of bast fibre from the stem of the plant. The fibre may be used as a substitute for jute in making burlap. The red calyces of the plant are used as food colourings and dyes. Today, roselle is attracting the attention of food and beverage manufacturers and pharmaceutical concerns who feel it may have exploitable possibilities as a natural food product for herbal medicine and as a colorant to replace some synthetic dyes.

#### *Nutritive value*

The nutritional analysis of roselle plant by proximate method (Luvonga et al. 2010) found the

carbohydrate content (68.7%) was highest followed by crude fibre (14.6%) and ash content (12.2%) and others. The plant is also found to be rich in minerals especially potassium and magnesium. Vitamins (ascorbic acid, niacin and pyridoxine) were also present in appreciable amounts. Various workers (Nnam and Onyeke 2003; Ojokoh 2006; Falade et al. 2005; Adanlawo and Ajibade 2006) reported variable content suggesting that the type of soil influences its ash and mineral content causing variations within the same species (Carvajal et al. 2012). It has long been used in herbal tea to treat hypertension, pyrexia and liver damage although the pharmaceutical components are poorly defined (Hou et al. 2005). Nutritional studies have indicated that low consumption of fruits and vegetables is consistently related to an increased incidence of cancer (Choi and Mason, 2000) reflecting dietary habits. The component in fruits and vegetables like polyphenol and anthocyanin may be responsible for the reduced risk of cancer (Weisburger and Chung 2002; Mei et al. 2005; Lin et al. 1999; Wang et al. 2003; Gao et al. 2002; Briviba et al. 2001). Plants have the capacity of producing secondary metabolites like proteins, steroids, alkaloids, etc (Sharaniah et al. 2013) that will enhance its nutritive value.

Animal Health Division, ICAR-RC for NEH Region, Umiam – 793103. Meghalaya

\*Corresponding author's Email: akulepuro@rediffmail.com

### *Antimicrobial properties*

Roselle is widely used for the treatment of diseases. Olaleye (2007) used the aqueous-methanolic extract of roselle to investigate its phytochemical constituents, antimicrobial activity and cytotoxicity, and reported that the extract contained cardiac glycosides, flavonoids, saponins and alkaloids. It exhibited antibacterial activities against *Staphylococcus aureus*, *Bacillus stearotherophilus*, *Micrococcus luteus*, *Serratia marseilles*, *Clostridium sporogenes*, *Escherichia coli*, *Klebsiella pneumoniae*, *Bacillus cereus*, *Pseudomonas fluorescens*. The results support the use of this plant in the treatment of diseases like abscesses, bilious conditions, cancer and coughs in traditional medicine, and also suggest the possibility of isolating antibacterial and anticancer agents while the antimicrobial activity on *Escherichia coli* O157:H7, *Salmonella enterica* and *Listeria monocytogenes* isolates from food, veterinary, and clinical samples by Fullerton (2011) indicated that roselle extract was effective and suggest the application of extracts as potential antimicrobials in foods. The antibacterial effects of roselle calyx aqueous and ethanol extracts and protocatechuic acid against food spoilage bacteria *Salmonella Typhimurium* DT104, *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Staphylococcus aureus* and *Bacillus cereus* were examined by Chau et al. (2008) and shown the inhibitory activity in dose-dependent manner against test bacteria in ground beef and apple juice and suggested that it might be potent agents as food additives to prevent contamination from these bacteria.

### *Antioxidant properties*

Roselle - Hibiscus anthocyanins (HAs) which are a group of natural pigments existing in the dried calyx exhibited antioxidant activity and liver protection. HA antioxidant bioactivity in rat primary hepatocytes and hepatotoxicity was studied by Wang et al. (2000). The results demonstrated that HAs, at the concentrations of 0.10 and 0.20mg/ml, significantly decreased the leakage of lactate dehydrogenase and the formation of malondialdehyde and significantly lowered the serum levels of hepatic enzyme markers (alanine and aspartate aminotransferase) and reduced oxidative liver damage. The histopathological evaluation of the liver revealed that roselle pigments reduced the incidence of liver lesions including

inflammatory leucocyte infiltration, and necrosis induced by tert-butyl hydroperoxide (t-BHP) in rats. An antioxidative activity was also reported in cancerous cell lines (Akim et al. 2011). In animal models (McKay et al. 2011), extracts of its calyces have demonstrated hypocholesterolemic and antihypertensive properties irrespective of age, gender or dietary supplement used. The antioxidant potential of three fractions of the ethanol crude extract (HS-C: chloroform-soluble fraction; HS-E: ethylacetate soluble fraction; HS-R: residual fraction) obtained from the dried flowers were evaluated by Tseng et al. (1997) for their capacity to quench free radicals and inhibiting xanthine oxidase (XO) activity. HS-E showed the greatest capacity of scavenging free radical, and HS-C showed the strongest inhibitory effect on XO activity. Furthermore, antioxidant bioactivities of these crude extracts were investigated in rat primary hepatocytes. All fractions were found to inhibit significantly the unscheduled DNA synthesis (UDS). These results indicated that the dried flower extracts (HS-C and HS-E) protect rat hepatocytes from t-BHP-induced cytotoxicity and genotoxicity. The study on hepatoprotective and antioxidant effects on the carbon tetrachloride (CCl<sub>4</sub>)-induced hepatocyte damage in fish by Yin et al. (2011) provided evidence of potential use as a medicine for curing liver diseases in aquaculture as roselle extract significantly elevated levels of lactate dehydrogenase (LDH), glutamate oxalate transaminase (GOT), glutamate pyruvate transaminase (GPT), and malondialdehyde (MDA) and significantly reduced levels of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px).

### *Anticancerous properties*

The antiproliferative activities of roselle juice were evaluated by Akim et al. (2011) using different cell lines like ovarian (Caov-3), breast (MCF-7, MDA-MB-231) and cervical (HeLa) cancer cell lines and found that it exhibited the strongest antiproliferative potency towards the MCF-7 cancer cells. The effects on human cancer cells (HL-60) studied by Chang et al. (2005) using roselle-anthocyanins (HA) showed apoptosis of cells in a dose- and time-dependent manner. It also revealed increased phosphorylation in p38, c-Jun and cytochrome c release, and expression of tBid, Fas, and FasL genes indicating that it could be developed

as chemopreventive agents. However, Hou et al. (2005) reported the apoptosis of leukemia cells induced by anthocyanin is through reactive oxygen species mediated mitochondrial pathways. Protocatechuic acid (PCA), a phenolic compound isolated from the dried flower, was found to inhibit the survival of human promyelocytic leukemia (HL-60) in a concentration and time dependent manner (Tseng et al. 1997), and apoptosis is induced via reduction of retinoblastoma phosphorylation and down regulation of Bcl-2 protein expression (Tseng et al. 2000). The study revealed that cells underwent intranucleosomal DNA fragmentation and morphological changes characteristics of apoptosis while the action against gastric carcinoma cells by inducing apoptosis is through JNK/MAPK signaling pathways (Lin et al. 2007). The methanolic extract of roselle on seven cancer lines (Lin et al. 2005) implied the AGS cancer cells being most susceptible in concentration-dependant form affecting both the intrinsic and extrinsic apoptotic routes.

#### *Effect on lipid metabolism*

The effect of roselle on lipid profile, creatinine and serum electrolytes has been studied by Abbas et al. (2011) in hypertensive patients and reported the upward trend of total cholesterol and high density lipid (HDL) which is significant since HDL-Cholesterol is a protective factor for coronary heart diseases. Kirdpon (1994) evaluated the changes of urine in normal patients after consuming roselle juice in different concentrations and durations which may help the treatment and prevention of renal stone disease, and reported a decrease of creatinine, uric acid, citrate, tartrate, calcium, sodium, potassium and phosphate but not oxalate in urinary excretion.

#### *Antihypertensive effect*

The consequence of hypertension is implicated in the development of cerebrovascular diseases, cardiac ischemia as well as cardiac and renal failure, and is now considered a global health problem. The studies on the efficacy of aqueous extract in hypertensive human (Faraji et al. 1999) showed significant reduced pressure difference in both systolic and diastolic compared to control group, while McKay et al. (2010) found the decrease in systolic pressure significant, the diastolic pressure remained unchanged. Studies were also conducted

in rats (Onyenekwe et al. 1999; Odigie et al. 2003; Ajay et al. 2007), and findings support the popular belief that roselle extract contains antihypertensive constituents. The anthocyanins extract investigated for its therapeutic efficacy, safety and tolerability along with antihypertensive drug captopril (Herrera-Arellano et al. 2004), lisinopril (Herrera-Arellano et al. 2007) in humans found the results comparable and suggest the synergistic mechanism of diuretic and ACE inhibition results in exerting hypotensive effects.

#### *Effect on domestic animal studies*

Few studies in animals have been reported. Roselle extract as acidifiers has been shown by Aphirakchatsakun et al. (2008) in post weaning pig with the ability to increase trypsin activity, fat digestibility and improve feed conversion ratio (FCR). In poultry, the effect of roselle calyx in layer diets on egg production performance, egg quality and Thiobarbituric acid reactive substances (TBARS) value in plasma and yolk was studied by Piyaphon et al. (2011) to check the lipid peroxidation as a result of degradation of fats. Storage time of extract was found to be an important factor to decrease egg quality and increase TBARS value in yolk.

#### *Future approach*

Among the properties reported to date, its effect on lipid metabolism, antihypertensive action and apoptosis are largely studied. Some studies on its antimicrobial effects were also documented. Therefore, with many enriching chemical-biological knowledge from animal and human models using plant extracts, future studies with greater scientific robustness in terms of standardization of dose for its effectiveness, safety and tolerability will permit the formulation of safe, effective therapeutic herbal formulations which can be used as an acceptable source for curing many health issues and restoring general health.

## REFERENCES

- Abbas M, Shirin M, Patricia K, Mohammad GK (2011). The effect of *Hibiscus Sabdariffa* on lipid profile, creatinine and serum electrolytes: A randomized clinical trial. ISRN Gastroenterol 2011: 1-4
- Adanlawo IG, Ajibade VA (2006). Nutritive value of the two varieties of roselle (*Hibiscus sabdariffa*) calyces soaked with wood ash. Pak J Nutr 5: 555-557

- Ajay M, Chai HJ, Mustafa AM (2007). Mechanisms of the anti-hypertensive effect of *Hibiscus sabdariffa* L. calyces. *J Ethnopharmacol* 109: 338-393
- Akim A, Lim CH, Asmah R, Zanainaul AZ (2011). Antioxidant and anti-proliferative activities of Roselle juice on Caov-3, MCF-7, MDA-MB-231 and Hela cancer cell lines. *Afri J Pharma Pharmacol* 5(7): 957-965
- Aphirakchatsakun W, Kris A, Suwanna K (2007). The effect of Roselle (*Hibiscus sabdariffa* Linn.) calyx as antioxidant and acidifier on growth performance in postweaning pigs. *Asian-Aust J Anim Sci* 21(4): 574-581
- Briviba K, Abrahamse SL, Pool-Zobel BL, Rechkemmer G (2001). Neurotensin-and EGF-induced metabolic activation of colon carcinoma cells is diminished by dietary flavonoid cyanidin but not by its glycosides. *Nutr Cancer* 41: 172-179
- Carvajal O, Maria D, Dremetriz B, Flores ZO, Margaret P, Jones H (2012). *Hibiscus sabdariffa* L., roselle calyx, from ethnobotany to pharmacology. *J Expt Pharmacol* 4: 25-39
- Chang YC, Huang HP, Hsu JD, Wang CJ (2005). *Hibiscus* anthocyanins rich extract-induced apoptotic cell death in human promyelocytic leukemia cells. *Toxicol Appl Pharm* 205: 201-212
- Chao C, Yin M (2008). Antibacterial Effects of Roselle Calyx Extracts and protocatechuic Acid in Ground Beef and Apple Juice. *Foodborne Pathogen Dis* 6(2): 201-206
- Choi SW, Mason JB (2000). Folate and carcinogenesis: an integrated scheme. *J Nutr* 130: 129-132
- Falade OS, Otemuyiwa IO, Oladipo A (2005). The chemical composition and membrane stability activity of some herbs used in local therapy for anemia. *J Ethnopharmacol* 102: 15-22
- Fullerton M, Khatiwada J, Johnson JU, David S, William LL (2011). Determination of antimicrobial activity of sorrel (*Hibiscus sabdariffa*) on *E.coli* O157:H7 isolated from food, veterinary and clinical samples. *J Med Food* 14(9): 950-956
- Gao X, Xu YX, Divine G, Janakiraman N, Chapman RA, Gautam SC (2002). Disparate in vitro and in vivo antileukemic effects of resveratrol, a natural polyphenolic compound found in grapes. *J Nutr* 132: 2076-2081
- Haji-Faraji MH, Haji-Tarkhani AH (1999). The effect of sour tea (*Hibiscus sabdariffa*) on essential hypertension. *J Ethnopharmacol* 65: 231-236
- Herrera-Arellano A, Flores-Romero S, Chávez-Soto MA (2004). Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild moderate hypertension: a controlled and randomized clinical trial. *Phytomed* 11: 375-382
- Herrera-Arellano A, Miranda-Sanchez J, Avila-Castro P, et al. (2007). Clinical effects produced by a standardized herbal medical product of *Hibiscus sabdariffa* on patients with hypertension. A randomized, double-blind, lisinopril-controlled clinical trial. *Planta Med* 73: 6-12
- Hou D, Tong X, Terahara N, Luo D, Fujii M (2005). Dephinidin 3-sambubioside, a *Hibiscus* anthocyanin, induces apoptosis in human leukemia cells through reactive oxygen species-mediated mitochondrial pathway. *Arch Biochem Biophys* 440: 101-109
- Kirdpon S (1994). Changes in urinary chemical composition in healthy volunteers after consuming roselle (*Hibiscus sabdariffa*) juice. *J Med Assoc Thai* 77(6): 314-21
- Lin H, Chen J, Kuo W, Wang C (2007). Chemopreventive properties of *Hibiscus sabdariffa* L. on human gastric carcinoma cells through apoptosis induction and JNK/p38 MAPK signaling activation. *Chemico-Biol Interact* 165: 59-75
- Lin H, Huang H, Huang C, Chen J, Wang C (2005). *Hibiscus* polyphenol rich extract induces apoptosis in human gastric carcinoma cells via p53 phosphorylation and p38 MAPK/FasL cascade pathway. *Mol Carcinog* 43: 86-99
- Lin JK, Liang YC, Lin-Shiau SY (1999). Cancer chemoprevention by tea polyphenols through mitotic signal transduction blockade. *Biochem Pharmacol* 58: 911-915
- Luvonga WAL, Njorge MS, Makokha A, Ngunjiri PW (2010). Chemical characterization of *Hibiscus sabdariffa* (Roselle) calyces and evaluation of its functional potential in the food industry. Proceeding of JKUAT Scientific and Industrial Conference. Kenya. pp 631-638
- Mckay DL, Chen CY, Saltzman E, Blumberg JB (2010). *Hibiscus sabdariffa* L. tea (tisane) lowers blood pressure in prehypertensive and mildly hypertensive adults. *J Nutr* 140: 298-303
- Mei Y, Wei D, Liu J (2005). Modulation effect of tea polyphenol toward *N*-methyl-*N*-nitro-*N*-nitrosoguanidine-induced precancerous gastric lesion in rats. *J Nutr Biochem* 16: 172-177
- Nnam NM, Onyike NG (2003). Chemical compositions of two varieties of sorrel (*Hibiscus sabdariffa* L.), calyces and the drinks made from them. *Plant Foods Hum Nutr* 58: 1-7
- Odigie IP, Ettarh RR, Adigun SA (2003). Chronic administration of aqueous extract of *Hibiscus sabdariffa* attenuates hypertension and reverses cardiac hypertrophy in 2K-1 hypertensive rats. *J Ethnopharmacol* 86:181-185
- Ojokoh AO (2006). Roselle (*Hibiscus sabdariffa*) calyx diet and histopathological changes in liver of albino rats. *Pak J Nutr* 5:110-113
- Olaleye MT (2007). Cytotoxicity and antibacterial activity of methanolic extract of *Hibiscus sabdariffa*. *J Med Plant Res* 1: 9-13
- Onyenekwe PC, Ajani EO, Ameh DA (1999). Antihypertensive effect of roselle (*Hibiscus sabdariffa*) calyx infusion in spontaneously hypertensive rats and a comparison of its toxicity with that in Wistar rats. *Cell Biochem Funct* 17: 199-206
- Piyaphon S, Kris A, Suwanna K (2011). Effect of Roselle (*Hibiscus sabdariffa* Linn.) calyx in laying hen diet on egg production performance, egg quality and TBARS value in plasma and yolk. *Thai J Vet Med* 41(3): 337-344
- Sharanaiah U, Shirin M, Mohammed A (2013). Antioxidant and antidiabetic activities of medicinal plants: a short review. *J Res Phytochem Pharmacol* 3(1): 40-53
- Tseng TH, Kao ES, Chu CY, Chou FP, Lin WHW, Wang CJ (1997). Protective effects of dried flower extracts of *Hibiscus sabdariffa* L. against oxidative stress in rat primary hepatocytes. *Food Chem Toxicol* 35: 1159-1164
- Tseng T, Kao T, Chu C, Chou F, Lin W, Wang C (2000). Induction of apoptosis by *Hibiscus* protocatechuic acid in human leukemia cells via reduction of retinoblastoma (RB) phosphorylation and Bcl-2 expression. *Biochem Pharmacol* 60: 307-315
- Wang CJ, Wang JM, Lin WL, Chu CY, Chou FP, Tseng TH (2000). Protective effects of *Hibiscus* anthocyanins against

- tert-butyl hydroperoxide-induced hepatic toxicity in rats. Food Chem Toxicol 38: 411-416
- Wang S, DeGroot VL, Clinton SK (2003). Tomato and soy polyphenols reduce insulin-like growth factor-I-stimulated rat prostate cancer cell proliferation and apoptotic resistance *in vitro* via inhibition of intracellular signaling pathways involving tyrosine kinase. J Nutr 133: 2367-2376
- Weisburger JH, Chung FL (2002). Mechanisms of chronic disease causation by nutritional factors and tobacco products and their prevention by tea polyphenols. Food Chem Toxicol 40: 1145-1154
- Yin G, Cao I, Xu P, Jenny G, Nakao M (2011). Hepatoprotective and antioxidant effects of *Hibiscus sabdariffa* extract against carbon tetrachloride-induced hepatocyte damage in *Cyprinus carpio*. In vitro Cellular Develop Biol Animal 47: 10-15